

CHECKLIST OF THE ORIBATID MITES OF THE NETHERLANDS (ACARI: ORIBATIDA)

Henk Siepel, Andrei Zaitsev & Matty Berg

More than fifty years ago Van der Hammen published the last checklist of oribatid mites (or moss mites) for the Netherlands. Since then the species number has almost doubled to 318 species, of which 100 are presented here for the first time. Brief data on occurrence and nomenclature are provided for each species. A short list of fossil species known from the Netherlands is presented separately.

INTRODUCTION

Oribatid mites (fig. 1) form an extremely diverse group of soil micro-arthropods with about 11.000 known species world-wide (Walter & Proctor 1999). They belong to the Chelicerata (spiders and alike) and are placed in the suborder Oribatida s.l. of the order Sarcoptiformes. Coddington et al. (2004) place the Oribatida s.s. in the order Sarcoptiformes parallel to the Astigmata. In this list the Astigmata are not included, but more primitive oribatids are, therefore we call it Oribatida s.l. in the traditional opinion. In their natural habitats oribatids may reach a density of ten to a

hundred thousand individuals per square meter. Up to sixty species may be found in one soil sample with a diameter of ten centimeters. Due to their small size (oribatids are usually less than 1 mm long) and high diversity identification to the species level is difficult. Moreover, the taxonomic status of many species has not always been clear.

Oribatid mites form an important part of detritus food webs. Most species feed on fungi and litter and have a large influence on the decomposition of dead organic matter. A number of species climb into the vegetation and trees and some




Figure 1. *Synchthonius crenulatus*.
Source: Van der Hammen (1952).
Figuur 1. *Synchthonius crenulatus*.
Bron: Van der Hammen (1952).

of them even have an arboricol way of life. Many predators are feeding on oribatid mites, such as gamasid mites, centipedes, Coleoptera such as Ptiliidae, Scydmaenidae and Pselaphidae and even salamanders (Krivolutsky 1995). Knowledge of the composition of oribatid mite communities is very relevant because of the significant role of this group in the soil ecosystem.

RESEARCH IN THE NETHERLANDS

The history of investigations on the oribatid mite fauna in the Netherlands dates back to the 19th century. The first checklist with about 60 species was published by Oudemans (1896). A revised edition of the list was published by Oudemans (1902). Van der Hammen (1952) further explored the oribatid fauna. He described several new species in the family Malaconothridae. His interest in oribatids enlarged the revised list to 162 species. More than half a century has passed since then. From the 1950s on much research was devoted to the ecology of oribatid mites, mainly by soil ecologists. This activity has resulted in many new records for the Dutch fauna which to date have not yet been published.

INTRODUCTION TO THE SPECIES LIST

The authors have revised the checklist published by Van der Hammen (1952). This has resulted in a total of 318 species, of which 100 are presented for the first time. Species were included only if at least one voucher specimen was available from the Netherlands. The few exceptions to this rule, when the specimen was lost, are discussed in the notes. The voucher specimens are located in the collections of the authors, of the Natural History Museums of Leiden and Amsterdam, of the research institute Alterra in Wageningen, or in the collection of the Vrije Universiteit in Amsterdam. The majority of literature and museum records were compiled by the first author. The resulting checklist was complemented by the other authors with 17 new species for the Netherlands collected during research on oribatid mite diversity in

forests, conducted during spring and summer 2000 (Zaitsev & Berg 2001). Workers of Naturalis and the European Invertebrate Survey - the Netherlands put much effort in digitalizing the data of RMNH, which contributed significantly to the total number of data for this checklist. Species not mentioned before in the literature are indicated with an asterisk (*).

The following species were not included into the list:

Nothrus borussicus Sellnick, 1928 (family Nothridae) was reported by Krivolutsky (pers. comm.) to exist in the Noord-Hollands Duinreservaat in North-Holland near IJmuiden and in 'het Amsterdamse Bos' in Amsterdam. Unfortunately a slide with the specimen of this species was not provided. Also older records from literature exist, but none of the specimens have been kept for examination. As confusion may have occurred with *N. anauniensis*, we remove the species from the checklist for the moment. An as yet undescribed species of the genus *Suctobelbella* (family Suctobelbidae) was found by D.A. Krivolutsky in 'het Amsterdamse Bos'. A so far unknown species of the genus *Brachychthonius* (family Brachychthoniidae) was collected by the second author near Bergen, Noord-Hollands Duinreservaat. This species resembles a species known only from Japan, *B. gracilis* (Chinone, 1974). The taxonomic status of the species is currently under investigation. An undescribed species of an unknown genus was found by the first author in old brooks (two records: Beekbergen en Nieuwkoop). The specimens probably belong to a species near the Parakalumnidae.

The systematics and nomenclature used in the checklist predominantly follows the work of Weigmann (2006); older reviews have been carefully checked (Balogh & Balogh 1992, Subias 2004). Our list was not checked extensively yet against the recent list of Fauna Europaea (Niedbala 2004).

We do not give any subdivisions below the

Oribatida to group the families. Current developments in phylogeny may change these groupings again on a short notice.

In the checklist species are followed by short notes on distribution data; habitat, locality, date and literature and collection references are given for the rare species only. Some important synonyms and misidentifications are provided. For the more common species reference has been made to the main collections in which specimens are present. The main source is a large database (and reference collection) at Alterra Green World Research Wageningen (ALT), furthermore the private collection of the first author (hs), collections at Naturalis Leiden (RMNH), Free University Amsterdam (VUA) and sources from literature.

SPECIES RECORDED AS SUBFOSSIL

Next to the list presented here, several species have been recorded from the Netherlands as subfossils. As the exoskeleton of oribatid mites is conserved very well in many soils (especially peat and clay soils), they can serve as indicators for ancient biotopes. Schelvis (1992) reconstructed the ancient biotopes of many archaeological excavations. The age of these subfossils goes as far as 40,000 years BP. It is obvious that in that period (during the age of the mammoths) the habitats in the Netherlands were quite distinct from the recent ones. Some of the oribatid mites, however, may persist in rare refugia till present. The following 11 species were recorded in the Netherlands as subfossil only by Schelvis (1992): *Epidamaeus bituberculatus* (Kulczynski, 1902), *E. tetricus* (Kulczynski, 1902), *Tectocephalus concavatus* Knölle, 1953, *T. translamellatus* Knölle 1954, *Ameronothrus lineatus* (Thörell, 1871), *Zygoribatula propinquus* (Oudemans, 1902), *Sphaerozetes orbicularis* (C.L. Koch, 1836), *Punctoribates hexagonus* Berlese, 1908, *Eupelops claviger* (Berlese, 1916), *Anachipteria dubia* Weigmann, 2001 and *Tegoribates latirostris* (C.L. Koch, 1844).

LIST OF SPECIES OF ORIBATID MITES OF THE NETHERLANDS

Class Arachnida

Subclass Acari

Order Sarcoptiformes

Suborder Oribatida s.l. (excl. Astigmata)

PALAEACARIDAE

Palaeacarus Trägårdh, 1932

1 *hystrinicu*s Trägårdh, 1932

Palaeacarus gallicus Grandjean, 1932

Few specimens sampled, probably quite common in low densities in old and undisturbed forests and hedgerows. Several records by Zaitsev & Berg (2001); Ughelen, 3.v.1994, hs.

APHELACARIDAE

Aphelacarus Grandjean, 1932

2* *acarinu*s (Berlese, 1910)

One record, several specimens, Academic hospital Leiden in culture with Psocoptera, 28.i.1981, RMNH.

BRACHYCHTHONIIDAE

Brachychthonius Berlese, 1910

3 *berlesei* Willmann, 1928

Brachychthonius brevis; Berlese 1910 [misidentification]
Several records Zaitsev & Berg (2001); Goeree, Springerborg, 11.III.1974, RMNH; in *Marchantia* at a steep slope of a stream Döttekro, Winterswijk, 9.IV.1993, hs.

4* *bimaculatus* Willmann, 1936

Brachychthonius helveticus Schweizer, 1956
In soil in an old grassland Bennekom (Bovenbuurt), 2.x.1989, hs.

5* *impressus* Moritz, 1976

Brachychthonius berlesei; Niedbala 1974 [misidentification]
In *Mnium undulatum*, Leudal, Haelen, 8.x.1989, hs.

- 6* **pius Moritz, 1976**
Brachychthonius berlesei; Perez-Inigo 1968
 [misidentification]
 Beech forest east of Nijmegen, 24.IX.1989, hs.
- Eobrachychthonius Jacot, 1936**
- 7 **mooseri (Van der Hammen, 1950)**
 Only one specimen from the type locality:
 Maastricht, St. Pietersberg, 6.V.1949, Van
 der Hammen (1952).
- 8 **oudemansi Van der Hammen, 1952**
Brachychthonius laetepictus; Willmann 1931
 [misidentification]
Eobrachychthonius argentinensis Hammer,
 1958
 One specimen from the type locality: Breda,
 Burgst, 1.V.1912, Van der Hammen (1952);
 Arnhem, 9.IX.1965, RMNH.
- Liochthonius Van der Hammen, 1959**
- 9 **alpestris (Forsslund, 1958)**
 Zeereep, Zoutelande, 8.III.2000, Zaitsev &
 Berg (2001).
- 10 **brevis (Michael, 1888)**
Brachychthonius perpusillus Berlese, 1910
 Several records by Zaitsev & Berg (2001); oak
 forest east of Nijmegen, 3.IX.1989, hs.
- 11 **evansi (Forsslund, 1958)**
Brachychthonius horridus; Evans 1952 [misid-
 entification]
Liochthonius forsslundi levis Chinone, 1974
 Zwilbroek, Eibergen, 16.III.1999, Zaitsev &
 Berg (2001); soil and litter in an old forest
 stand at Loenen, 3.V.1994, hs.
- 12* **furcillatus (Willmann, 1942)**
Brachychthonius ensifer Strenzke, 1951
 In moss in a larch stand, Nijmegen, 15.X.1989,
 hs.
- 13* **horridus (Sellnick, 1928)**
 Soil in an old hedgerow, Harskamp, 25.IX.1997,
 hs.
- 14 **hystrinicus (Forsslund, 1942)**
Brachychthonius ocellatus Hammer, 1952
 Several records by Zaitsev & Berg (2001).
- 15* **lapponicus (Trägårdh, 1910)**
Hypochthonius brevis var. *lapponicus*
 Trägårdh, 1910
 Several specimen in *Ammophila* litter, Meyen-
 del (ZH), 21.I.1974, RMNH.
- 16* **peduncularis (Strenzke, 1951)**
 In floating sods in peat swamp, Westbroekse
 Zodden, Westbroek, 29.VI.1992, hs.
- 17 **perfusorius Moritz, 1976**
Brachychthonius furcillatus; Strenzke 1951
 [misidentification]
 Spanderswoud, Hilversum, 9.VII.2000, Zaitsev
 & Berg (2001); pine litter at Ysselstein (Lim-
 burg), 28.III.1993, hs.
- 18 **sellnicki (Thor, 1930)**
Brachychthonius scalaris Forsslund, 1942
Brachychthonius nodosus Willmann, 1952
 Several records by Zaitsev & Berg (2001); for-
 est litter (among *Ficaria verna*), Bekkendelle,
 Winterswijk, 9.IV.1993, hs.
- 19* **simplex (Forsslund, 1942)**
 Imbos (GE), 12.XI.1985, ALT (specimens not
 stored).
- 20 **strenzkei (Forsslund, 1942)**
Brachychthonius sellnicki; Strenzke 1951
 [misidentification]
 Several records by Zaitsev & Berg (2001);
 in *Rhytidadelphus squarosus* on sand dunes,
 Horsmeertjes, Texel, 6.II.1993, hs.
- 21 **tuxeni (Forsslund, 1957)**
Brachychthonius hystrinicus var. *tuxeni*
 Forsslund, 1957
 Ulvenhoutse Bos, Nieuw Ginnen, 24.V.2000,
 Zaitsev & Berg (2001); litter in old forest
 stand, Ugcelen, 3.V.1994, hs.
- Neobrachychthonius Moritz, 1976**
- 22* **magnus Moritz, 1976**
Neobrachychthonius marginatus magnus
 Moritz, 1976
 Deelerwoud (GE), 20.XI.1984, ALT (specimens
 not stored).
- 23* **marginatus (Forsslund, 1942)**
 Deelerwoud (GE), 14.I.1988, ALT (specimens
 not stored).

- Poecilochthonius Balogh, 1943**
- 24 **italicus (Berlese, 1910)**
 Brachychthonius brevis (Michael, 1888) var.
italicus Berlese, 1910
 Specimens sampled from *Querco-Carpinetum*
 litter, Gronsveld, 14.IX.1950, Van der Hammen
 (1952).
- 25* **spiciger (Berlese, 1910)**
 Brachychthonius brevis (Michael, 1888) var.
spiciger Berlese, 1910
 Brachychthonius rapaporti Balogh &
 Mahunka, 1968
 One record from Zwilbroek, Eibergen
 9.IV.1993, hs.
- Sellnickochthonius Krivolutsky, 1964**
- 26 **cricoides (Weis-Fogh, 1948)**
 Several records by Zaitsev & Berg (2001).
- 27* **formosus (Cooreman, 1947)**
 In *Rhytidiaadelphus squarrosus* on sand dunes,
 Horsmeertjes, Texel, 6.II.1993, hs.
- 28* **honestus (Moritz, 1976)**
 Brachychthonius zelawaiensis; Evans 1952
 [misidentification]
 Brachychthonius zelawaiensis; Bulanova-
 Zachvatkina 1957 [misidentification]
 Beech forest east of Nijmegen, 24.IX.1989, hs.
- 29 **hungaricus (Balogh, 1943)**
 Sellnickochthonius rostratus Krivolutsky,
 1975
 One record Amsterdamse Bos, Amsterdam,
 15.VIII.2000, Zaitsev & Berg (2001); in
Rhytidiaadelphus squarrosus on sand dunes,
 Horsmeertjes, Texel, 6.II.1993, hs.
- 30 **immaculatus (Forsslund, 1942)**
 Brachychthonius semiornatus Evans, 1952
 Brachychthonius arcticus Hammer, 1952
 Brachychthonius striatus Willmann, 1956
 Brachychthonius gyperi Schweizer, 1963
 Brachychthonius obscurus Krivolutsky, 1966
 Heemskerk, 12.IV.2000, Zaitsev & Berg (2001);
 Goeree, Springergors, 6.VI.1975, RMNH (sub.
B. semiornatus).
- 31* **subcricoides (Balogh & Mahunka, 1979)**
 Hoge Veluwe, Heiweg, 20.XI.1984, ALT (speci-
 mens not stored).
- 32 **suecicus (Forsslund, 1942)**
 Brachychthonius jugatus var. *suecica*
 Forsslund, 1942
 Brachychthonius jugatus Niedbala, 1972
 Few localities, probably more widespread,
 Hoenderloo, Hoge Veluwe, 24.X.1942, Van der
 Hammen (1952).
- 33 **zelawaiensis (Sellnick, 1928)**
 Several records by Zaitsev & Berg (2001);
 old forest stand Ugchelense Bos, Ugchelen,
 3.V.1994, hs.
- Synchthonius Van der Hammen, 1952**
- 34 **crenulatus (Jacot, 1938)**
 Synchthonius boschmai Van der Hammen,
 1952
 Specimens from litter of a cultivated *Larix*
 with *Alnus* forest, Eext, 11.V.1943, Van der
 Hammen (1952); Molenveld, Saasveld,
 26.IV.2000, Zaitsev & Berg (2001).
- Verachthonius Moritz, 1976**
- 35* **congruus Moritz, 1976**
 Arnhemse Heide, 14.I.1988, ALT (specimens not
 stored).
- 36* **laticeps (Strenzke, 1951)**
 Deelerwoud (GE) and Imbos in litter bags,
 14.IV.1986, ALT (specimens not stored).
- COSMOCHTHONIIDAE**
- Cosmochthonius Berlese, 1910**
- 37* **lanatus (Michael, 1885)**
 Cosmochthonius domesticus Grandjean, 1947
 In *Thuja* litter, Nieuw Dennenlust,
 Beekbergen, 17.VII.1989, hs.
- HAPLOCHTHONIIDAE**
- Haplochthonius Willmann, 1930**
- 38 **simplex Willmann, 1930**
 One old record from the Netherlands but with
 some uncertainty concerning its precise locali-
 ty, sampled by Oudemans (Van der Hammen
 1952) and one more recent: Texel, polder Het
 Noorden, 14.VI.1972, RMNH.

ATOPOCHTHONIIDAE

Atopochthonius Grandjean, 1948

39* *artiodactylus* Grandjean, 1948

Litter and soil in an old hedgerow, Nunspeet,
25.IX.1997, HS.

HYPOCHTHONIIDAE

Hypochthonius C.L. Koch, 1836

40 *luteus* Oudemans, 1917

Valkenburg (type locality), Oudemans in Van der Hammen (1952); Stavenisse (Tholen) and Vlissingen, 3.XI.1974, RMNH; among litter with *Anemone nemorosa*, St. Pietersberg, Maastricht, 21.IV.1994, HS.

41 *rufulus* C.L. Koch, 1836

Several records by Van der Hammen (1952) and Zaitsev & Berg (2001); Hackfort, Vorden, 1959, Minderman (1981) and RMNH; slope of a ditch in *Mnium hornum*, Beekbergerwoud, Beekbergen, 14.VII.1991, HS.

ENIOCHTHONIIDAE

Eniochthonius Grandjean, 1933

42 *minutissimus* (Berlese, 1904)

Hypochthonius pallidulus Berlese, 1896
(misidentified type specimen)

Eniochthonius grandjeani Van der

Hammen, 1952

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; slope of a ditch in *Mnium hornum*, Beekbergerwoud, Beekbergen, 14.VII.1991, HS.

EULOHMANNIIDAE

Eulohmannia Berlese, 1910

43 *ribagai* Berlese, 1910

Arthronothrus biunguiculatus Trägårdh, 1910
Hackfort, Vorden, 1959, Minderman (1981); Arnhem, 18.VI.1965; Winterswijk, Willink Weuste, 1.VII.1975; Vijlenerbos (LI), 29.V.1974; Plasmolen, 31.X.1968; Epen, 6.V.1972;

Westerbroek (Gr), 20.VI.1984, Heerenveen, 18.VI.1979; Schoorl, 21.VI.1983, RMNH; records from old forest stands and hedgerows, Epe, Vemde, 22.IX.1997, HS.

EPILOHMANNIIDAE

Epilohmannia Berlese, 1910

44 *cylindrica* (Berlese, 1904)

Lesseria szanisloii Oudemans, 1917

Epilohmannia verrucosa Jacot, 1934

Two records: Nolledijk, Vlissingen, 20.II.1973; Koudekerke inlagen, Westerschouwen, 3.XI.1974 (Polderman & de Vries 1978) sub *E. szanisló*.

PHTHIRACARIDAE

Archiphthiracarus Balogh & Mahunka, 1979

45* *anonymus* (Grandjean, 1934)

Record from a birchbrook, Zwilbroek, Eibergen, 9.IV.1993, HS.

Atropacarus Ewing, 1917

46 *striculus* (C.L. Koch, 1836)

Steganacarus diaphanus Jacot, 1930

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; sampled from pine litter at Ysselstein (LI), 28.III.1993, HS.

Hoplothpiracarus Jacot, 1933

47 *illinoiensis* (Ewing, 1909)

Hoploderma pavidum; Van der Hammen 1963 [misidentification]

Hoplothpiracarus vanderhammeni Niedbala, 1991

Phthiracarus pavidus (Berlese, 1913)

Manteling van Walcheren, Domburg, 8.III.2000, Zaitsev & Berg (2001); Wijster, 9.VIII.1962, in Sphagnum near mere, RMNH.

Phthiracarus Perty, 1841

48 *crenophilus* Willmann, 1951

Spanderswoud, Hilversum, 9.VII.2000;

Agelerbroek, Ootmarsum, 26.IV.2000, Zaitsev & Berg (2001).

- 49 **crinitus** (C.L. Koch, 1841)
Uiterwaarden Waardenburg, 28.VIII.2000,
Zaitsev & Berg (2001).
- 50 **ferrugineus** (C.L. Koch, 1841)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; litter
between *Equisetum hyemale*, Bekkendelle,
Winterswijk, 9.IV.1993, HS.
- 51 **globosus** (C.L. Koch, 1841)
Several records by Van der Hammen (1952)
and RMNH; litter and moss near the stream
Leudal, Haelen, 8.X.1989, HS.
- 52 **laevigatus** (C.L. Koch, 1841)
Phthiracarus nitens (Nicolet, 1855)
Phthiracarus parabotrichus Feider & Suciuc,
1975
Phthiracarus piger (Scopoli, 1763)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; birchbrook
in peatland, Nieuwkoop, 2.VII.1992; litter of
conifers and broadleaved trees, Schaarsbergen,
Arnhem, 19.VII.1989, HS.
- 53 **longulus** (C.L. Koch, 1841)
Phthiracarus borealis; Willmann 1931 [misiden-tification]
Phthiracarus flexisetosus Parry, 1979
Phthiracarus hortobagyensis Mahunka, 1982
Phthiracarus mundus Niedbala, 1983
Phthiracarus setosellus Jacot, 1928
Phthiracarus tardus Forsslund, 1956
Moosbeek, Tubbergen, 26.IV.2000, Zaitsev &
Berg (2001).
- 54 **testudineus** (C.L. Koch, 1841)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; in moss
on *Prunus avium*, Savelbos, Gronsveld,
21.IV.1994, HS.
- Steganacarus Ewing, 1917**
- 55 **anomalus** (Berlese, 1883)
Steganacarus karamani Tarman, 1962
Several records by Zaitsev & Berg (2001);
among litter with *Anemone nemorosa*,
St. Pietersberg, Maastricht, 21.IV.1994, HS.
- 56* **applicatus** (Sellnick, 1920)
Phthiracarus berlesei Oudemans, 1915
(preocc.)
Mantinge (DR) 28.IX.1971, Bunderbos (LI)
28.V.1974, RMNH; litter of conifers and
broadleaved trees, Schaarsbergen, Arnhem,
19.VII.1989, HS.
- 57 **magnus** (Nicolet, 1855)
Hackfort, Vorden, 1959, Minderman (1981);
several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; in *Mnium
undulatum* at a steep slope of a stream,
Döttekro, Winterswijk, 9.IV.1993, HS.
- 58* **michaeli** Bernini & Avanzati, 1987
In soil of an old hedgerow near Vemde, Epe,
22.IX.1997, HS.
- 59 **spinosus** (Sellnick, 1920)
Steganacarus punctulatus Sergenko, 1985
Several records by Van der Hammen (1952);
Norg (DR), 10.X.1972, RMNH; in *Mnium
undulatum* at a steep slope of a stream,
Döttekro, Winterswijk, 9.IV.1993, HS.

EUPHTHIRACARIDAE

Eupthiracarus Ewing, 1917

60* **cribarius** (Berlese, 1904)

Several records in RMNH from the provinces
Limburg and Gelderland; in moss in an oak
forest, Beetsterzwaag, 11.X.1989; decaying
wood, Heerenveen, 18.VI.1979; from moss on
fallen tree, Westerbroek (GR), 20.VI.1984, HS.

61 **monodactylus** (Willmann, 1919)

Eupthiracarus mixtus Mahunka, 1979
Bekkendelle, Winterswijk, 16.III.1999, Zaitsev
& Berg (2001); Arnhem, 28.IV.1965; Lankhorst
(Nijmegen), 20.V.1975; Mariahoop (Echt),
29.VI.1978; Haelen, 28.VI.1978; Broek in
Waterland, 5.VI.1984; Oldematen, Zwartewater
(OV), 10-11.VI.1985, RMNH; forest litter, Bekken-
delle, Winterswijk, 9.IV.1993, HS; all in litter.

- Microtritia Märkel, 1964**
- 62 **minima** (Berlese, 1904)
- Tritia (Pseudotritia) minuta Willmann, 1919
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; litter in old
forest stand, Ugchelen, 3.v.1994, hs.
- Rhysotritia Märkel & Meyer, 1959**
- 63 **ardua** (C.L. Koch, 1841)
- Tritia lentula Berlese, 1887
- Phthiracarus canestrinii Michael, 1898
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; birchbrook,
Zwolbroek, Eibergen, 9.iv.1993, hs.
- 64 **duplicata** (Grandjean, 1953)
- Several records by Zaitsev & Berg (2001) and
RMNH; litter in old forest stand, Ugchelen,
3.v.1994, hs.
- MALACONOTHRIDAE**
- Malaconothrus Berlese, 1904**
- It is possible that all species described by
Van der Hammen, 1952 are varieties of
M. monodactylus (Michael, 1888); the species
has a thelytokous reproduction and forma-
tion of distinct clones is to be expected.
However, ecological preferences may occur
and therefore the separate species are listed
for the moment.
- 65 **gracilis** Van der Hammen, 1952
- Few records from alder marsh and the source
of a brook, Ootmarsum, 22.iv.1949; De Lutte,
21.iv.1949, Van der Hammen (1952); moss on
tree trunk, Ittervoort, Vijverbroek, 28.III.1993,
hs.
- 66 **mollisetosus** Hammer, 1952
- Several records by Zaitsev & Berg (2001).
- 67* **monodactylus** (Michael, 1888)
- Malaconothrus egregius Berlese, 1910
Birchbrook, Zwolbroek, Eibergen, 9.iv.1993, hs.
- 68 **processus** Van der Hammen, 1952
- Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001); in *Mnium* on
border of a ditch, Beekbergen, Elsbosweg,
14.VII.1991, hs.
- 69 **punctulatus** Van der Hammen, 1952
- Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001); in moss on soil,
Mook, Helkuil, 28.VIII.1989, hs.
- Trimalaconothrus Berlese, 1916**
- 70* **angulatus** Willmann, 1931
- Veenendaal, Blauwe Hel, peat swamp,
31.VIII.1989, hs.
- 71* **glaber** (Michael, 1888)
- Terschelling, Dellewal, 22.IV.1975; Wieringen,
Normerdijk, 12.XII.1975, RMNH.
- 72 **grandis** Van der Hammen, 1952
- Record from the type locality, Rockanje,
17.IX.1948, Van der Hammen (1952); Rockanje,
1.IX.1968, RMNH.
- 73 **maior** (Berlese, 1910)
- Malaconothrus novus Sellnick, 1921
Trimalaconothrus tardus; Buitendijk 1945
[misidentification]
Camisia monodactyla; Oudemans 1925 [mis-
identification]
Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001); Ameland,
29.VI.1972, RMNH (all sub *T. novus*); birch-
brook, Zwolbroek, Eibergen, 9.iv.1993, hs.
- TRHYPOCHTHONIIDAE**
- Trhypochthonius Berlese, 1904**
- 74 **cladonicola** (Willmann, 1919)
- Few records, in lichens in heathland, maybe
decreasing, Hoenderloo, 11.X.1948; Lattrop
22.IV.1949, Van der Hammen (1952); Ommen,
Lemelerberg, 19.VI.1973, RMNH.
- 75* **tectorum** (Berlese, 1896)
- Trhypochthonius nigricans; Jørgensen 1934
[misidentification]
Few records, but probably more widespread.
Ameland, 19.X.1975; Ubbergen, 19.XII.1977;
Meyendel (ZH), 21.I.1973; Rijnsburg, 7.XII.1954;
Utrecht, 13.V.1975; Schouwen Duiveland,
Oosterland, 6.X.1974; Vlissingen, 20.II.1973,
RMNH; Bergen (NH), date unknown; Boxtel,
in moss on roof tiles, 20.VII.1989, hs.

TRHYPOCHTHONIELLIDAE

Trhypochthoniellus Willmann, 1928

76 excavatus (Willmann, 1919)

Hoge Veluwe in *Sphagnum*, IJzeren Man, 11.X.1948; Van der Hammen (1952); in pond with *Sphagnum*, Beekbergen, Recenberg, 19.VII.1989, HS; Ommen, Beerzerveld, 20-21.VI.1973, RMNH.

NOTHRIDAE

Nothrus C.L. Koch, 1835

77 anauniensis Canestrini & Fanzago, 1876

Nothrus silvestris anauniensis Canestrini & Fanzago 1876
Camisia biciliatus; Oudemans 1902 [misidentification]
Nothrus biciliatus; Sellnick & Forsslund 1955 [misidentification]
Forest litter, Maastricht, St. Pietersberg, 6.V.1949; Van der Hammen (1952); Hackfort, Vorden, 1959; Minderman (1981); several records by Zaitsev & Berg (2001) and RMNH; birch brook, Nieuwkoop, 2.VII.1992; Maastricht, St. Pietersberg, 21.IV.1994, HS.

78 palustris C.L. Koch, 1839

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Rhytidia-delphus squarosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, HS.

79* pratensis Sellnick, 1929

In marshes in *Sphagnum*, Spier, 23.VII.1949; Hoge Veluwe, 11.X.1948; Heerenveen, 18.VI.1979; Helumer Oldeferd, 19.VI.1979, Losser, 1.VI.1990; Ommen, 19-21.VI.1973; Vriezenveen, 28.VIII.1979; Winterswijk, Korenburgrerveen, 2.VII.1975; Putten, 20.IX.1969; Wijster, 9.VIII.1962; Delft, 21.V.1972; Bemelen, 15.X.1952, RMNH.

80 silvestris Nicolet, 1855

Nothrus anauniensis; Berlese 1885 [misidentification]

Angelia anauniensis; Lombardini 1936 [misidentification]

Several records by Van der Hammen (1952), Zaitsev & Berg (2001), HS, ALT and RMNH.

CAMISIIDAE

Camisia von Heyden, 1826

81 biurus (C.L. Koch, 1839)

Camisia exuvialis Grandjean, 1939

Several records by Van der Hammen (1952) and RMNH; litter around *Juniperus* shrubs Otterlo 6.IX.1989, HS.

82 biverrucata (C.L. Koch, 1839)

Nothrus horridus; Nicolet 1855 [misidentification]

Camisia fischeri Oudemans, 1900

Few records on dry vegetations, Meyendel, 24.III.1949; Van der Hammen (1952) and RMNH; Zwarte polder, Zeeuws Vlaanderen, 14.VII.1974, RMNH.

83* horrida (Hermann, 1804)

Nothrus mutilus C.L. Koch, 1839

Nothrus bistriatus C.L. Koch, 1839

Nothrus sinuatus C.L. Koch, 1839

Nothrus runcinatus C.L. Koch, 1839

Nothrus angulatus; Berlese 1883 [misidentification]

Nothrus rugulosus Banks, 1895

Ameland, 9.VI.1983; Weerselo 29.V.1990;

Wellerlooise Bos (IJ), 1977; Weert, 1976;

Haelen, 28.VI.1978; Hilvarenbeek, 15.VI.1977;

Goeree, Springergors, 6.VI.1975; Overflakkee,

Ooltgensplaat, 1975, RMNH; litter of *Thuja*,

Beekbergen, Nieuw Dennenlust, 11.VII.1989,

HS.

84 invenusta (Michael, 1888)

Terschelling, De Kom, 31.X.1972, RMNH; Moss on stone, Arnhem, Schaarsbergen, 5.VII.1990, HS.

85 segnis (Hermann, 1804)

Nothrus bicarinatus C.L. Koch, 1839

Nothrus rostratus C.L. Koch, 1839

Nothrus ventricosus C.L. Koch, 1839

Nothrus excisus Banks, 1895

Several records by Van der Hammen (1952), RMNH, Zaitsev & Berg (2001) and Noordijk & Berg (2001); among conifer litter, Arnhem, Schaarsbergen, 4.VII.1989, HS.

- 86 **spinifer** (C.L. Koch, 1835)
Nothrus taurinus Banks, 1906
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.
- Heminothrus Berlese, 1913**
- 87 **targionii** (Berlese, 1885)
In forests, Maastricht, St. Pietersberg, 6.v.1949,
Van der Hammen (1952); litter between
Equisetum hyemale, Bekkendelle, Winterswijk,
9.IV.1993, HS.
- Platynothrus Berlese, 1913**
- 88 **peltifer** (C.L. Koch, 1839)
Nothrus palliatus C.L. Koch, 1839
Nothrus bistriatus; Nicolet 1855 [misidentification]
Nothrus cirrosus Canestrini & Fanzago
1876
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT, RMNH.
- 89* **thori** (Berlese, 1904)
Hasselt, II.VI.1985, RMNH; in moss in a peat swamp, Veenendaal, De Hel, 7.IX.1989, HS.
- NANHERMANNIIDAE**
- Nanhermannia Berlese, 1913**
- 90 **comitalis** Berlese, 1916
Nanhermannia ussurica Golosova, 1969
Specimens sampled from wet hayfields and moors, Maasland, 2.IV.1949, Van der Hammen (1952); several records from Zaitsev & Berg (2001) and RMNH.
- 91 **coronata** Berlese, 1913
Nanhermannia nana; Willmann 1931 [misidentification]
Several records by Zaitsev & Berg (2001) and RMNH; birchbrook, Zwilbroek, Eibergen, 9.IV.1993, HS.
- 92 **elegantula** Berlese, 1913
Nanhermannia areolata Strenzke, 1953
Several records by Van der Hammen (1952); oak litter, Hoge Veluwe, Franse Berg, 7.VIII.1989, HS.
- 93 **nana** (Nicolet, 1855)
Nanhermannia elegantula; Sellnick 1928
[misidentification]
Nanhermannia dorsalis elegantula Jacot, 1937
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, HS.
- 94 **pectinata** Strenzke, 1953
Agelerbroek, Ootmarsum, 26.IV.2000, Zaitsev & Berg (2001); Ameland, Eendekooi, 4.III.1972, RMNH; in *Marchantia* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, HS.
- 95 **sellnicki** Forsslund, 1958
Ulvenhoutshe Bos, Nieuw Ginniken, 24.V.2000, Zaitsev & Berg (2001); old hedgerow, Stroe, Walweg, 25.IX.1997, HS.
- HERMANNIIDAE**
- Hermannia Nicolet, 1855**
- 96 **gibba** (C.L. Koch, 1840)
Angelia piceus Berlese, 1885
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) & RMNH; moss on dead *Prunus*, Rijkholt, Savelbos, 21.IV.1994, HS.
- 97 **nodosa** Michael, 1888
Nothrus scabrus L. Koch, 1879 (partim)
Lohmannia scabra (L. Koch, 1879)
Records from the seashore from Oudemans (Putten, Zeeburg) and Geijskes (Griend) in Van der Hammen (1952); Ameland, 4.III.1972; 8.VI.1972, Holwerd; 16.X.1975, Gaasterland; 23.III.1973, Sas van Goes, 20.II.1973 and 9.XI.1972; Stavenisse (Tholen), 3.XI.1974; Rilland, 28.VIII.1975; Dieden, (NB), V.1975, RMNH; litter on the bank, Texel, Mokbaai, 6.II.1993, HS.
- 98* **subglabra** Berlese, 1910
Nothrus scabrus L. Koch, 1879 (partim)
Vast amount of records from the survey of P.G.J. Polderman, all seashore data at RMNH.

HERMANNIELLIDAE

- Hermannella Berlese, 1908**
99* dolosa Grandjean, 1931
 Litter between *Equisetum hyemale*,
 Bekkendelle, Winterswijk, 9.IV.1993, HS.
100 granulata (Nicolet, 1855)
 Several records by Van der Hammen (1952),
 Zaitsev & Berg (2001) and RMNH; Hackfort,
 Vorden, 1959, Minderman (1981); in moss,
 Winterswijk, Willink Weuste, 9.IV.1993, HS.
101 punctulata Berlese, 1908
Nothrus piceus C.L. Koch, 1839 (probably
 not *Hermannella*)
 Several records by Van der Hammen (1952);
 Hackfort, Vorden, 1959, Minderman (1981).
102* septentrionalis Berlese, 1910
Hermannella punctulata septentrionalis
 Berlese, 1910
 Litter under *Anemone nemorosa*, Maastricht,
 St. Pietersberg, 21.IV.1994, HS.

NEOLIODIDAE

- Poroliodes Grandjean, 1934**
103* farinosus (C.L. Koch, 1840)
Nothrus theleproctus Michael, 1888
 Vorden, Hackfort, 11.I.1962, Oldematen (ov),
 11.VI.1985, RMNH.

GYMNODAMAEIDAE

- Gymnodamaeus Kulczynski, 1902**
104* bicostatus (C.L. Koch, 1836)
 In epiphytes, St. Pietersberg (LI), 14.X.1952,
 RMNH.

LICNODAMAEIDAE

- Licnodamaeus Grandjean, 1931**
105 pulcherrimus (Paoli, 1908)
 Dry litter in sand dunes, Meyendel, 22.I.1949,
 22.III.1949, 1.IV.1949, Van der Hammen (1952);
 Meyendel, 21.I.1974, RMNH; Castricum, Vogel-
 duin, 24.X.2007; Wijk aan Zee, Noorderduin
 en De Rellen, 24.X.2007, ALT.

DAMAEIDAE

- Adamaeus Norton, 1977**
106 onustus (C.L. Koch, 1841)
Damaeus geniculatus C.L. Koch 1841
Belba geniculosa Oudemans, 1929
 Several records by Van der Hammen (1952),
 Zaitsev & Berg (2001) and RMNH; Hackfort,
 Vorden, 1959, Minderman (1981); in *Mnium*
 at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, HS.
Belba von Heyden, 1826
107 corynopus (Hermann, 1804)
Damaeus sufflexus Michael, 1885
 Several records by Van der Hammen (1952),
 Zaitsev & Berg (2001) and RMNH; litter un-
 der *Ficaria verna*, Winterswijk, Bekkendelle,
 9.IV.1993, HS.
108 bartosi Winkler, 1955
Belba pseudocorynopus Märkel & Meyer,
 1960
 One record, Bekkendelle, Winterswijk,
 16.III.1999, Zaitsev & Berg (2001).

Damaeobelba Sellnick, 1929

- 109 minutissima (Sellnick, 1920)**
 Several records by Van der Hammen (1952),
 Zaitsev & Berg (2001) and RMNH; Hackfort,
 Vorden, 1959, Minderman (1981); old forest
 stand in litter, Hoenderloose Bos, 3.V.1994, HS.

Damaeus C.L. Koch, 1835

- 110 auritus C.L. Koch, 1835**
Oribata gracilipes Kulczynski, 1902
 An old record from Denekamp by Oudemans
 and from Maastricht, St. Pietersberg, 6.V.1949
 and 22.IX.1949, Van der Hammen (1952);
 Ubbergen, 7.VI.1988; Groesbeek, De Bruuk,
 10.VI.1988; Waterop (LI), VI.1986; Huys te
 Linschoten, 4.VI.1985 (sub *D. gracilipes*),
 RMNH; in moss on soil, Mook, Helkuil,
 23.VIII.1989, HS.

- Epidamaeus Bulanova-Zachvatkina, 1957**
- 111 *glabriseta* (Willmann, 1930)
Belba verticillipes; Buitendijk 1945 [misidentification]
Records from Bergen and Nederweert (type specimens) sampled by Oudemans (Van der Hammen 1952).
- Hypodamaeus Bulanova-Zachvatkina, 1957**
- 112 *riparius* (Nicolet, 1855)
Oppia aurita Oudemans, 1900
Haarlem, Oudemans, Van der Hammen (1952); Heemskerk, 12.IV.2000, Zaitsev & Berg (2001).
- Kunstidamaeus Miko, 2006**
- 113* *tecticola* (Michael, 1888)
Roof of shed with reed, Ommen, Lemelerberg, 21.VI.1973, RMNH.
- Metabelba Grandjean, 1936**
- 114 *lanceolata* Van der Hammen, 1952
In *Sphagnum*, Lattrop (type locality), 22.IV.1949, Van der Hammen (1952).
- 115 *papillipes* (Nicolet, 1855)
Metabelba cremersi Van der Hammen, 1952
Damaeus michaelii Hull, 1915
Several records by Van der Hammen (1952) (sub *M. cremersi*); Winterswijk, Bekkendelle, RMNH; litter under *Ficaria verna*, Winterswijk, Bekkendelle, 9.IV.1993, HS.
- 116* *pulverosa* Strenzke, 1953
Oribata pulverulenta Kulcynski 1902 (misident. and suppressed, nec *Nothrus pulverulentus* C.L. Koch, 1839)
Meyendel (ZH), 30.V.1968 and 13.V.1971, Linschoten (UT), 4.VI.1985, RMNH (sub *M. pulverulenta*).
- 117 *rohdendorfi* Bulanova-Zachvatkina, 1965
Bekkendelle, Winterswijk, 16.III.1999, Zaitsev & Berg (2001) (sub *Belba* spec.).
- 118* *sphagni* Strenzke, 1950
Engbertsdijksveen (ov), 1974, RMNH;
quagmire, Westbroekse Zodden, 29.VI.1992,
HS.
- Paradamaeus Bulanova-Zachvatkina, 1957**
- 119 *clavipes* (Hermann, 1804)
Damaeus nodipes C.L. Koch, 1839
Acarus geniculatus Johnston, 1849
Several records by Van der Hammen (1952), Zaitsev & Berg (2001), HS, ALT and RMNH.
- Porobelba Grandjean, 1936**
- 120* *spinosus* (Sellnick, 1920)
Oldebroek (GE), 21.II.1961, RMNH; litter around *Juniperus* shrubs, Otterlo, 6.IX.1989, HS.
- Spatiodamaeus Bulanova-Zachvatkina, 1957**
- 121 *verticillipes* (Nicolet, 1855)
Belba torva Oudemans, 1897
Damaeus quadrihastatus Märkel & Meyer, 1960
Several records by Van der Hammen (1952), Zaitsev & Berg (2001), HS, ALT and RMNH.
- CEPHEIDAE**
- Cepheus C.L. Koch, 1835**
- 122 *cepheiformis* (Nicolet, 1855)
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Thuja* litter, Beekbergen, Nieuw Dennenlust, 11.VII.1989, HS.
- 123 *dentatus* (Michael, 1888)
Several records by Zaitsev & Berg (2001) and RMNH; beech forest east of Nijmegen, 3.IX.1989, HS.
- 124* *latus* C.L. Koch, 1835
In forest litter and dead wood, Vaals, Vijlenerbos and Schimperbos, 28.V.1982, RMNH.

Tritegeus Berlese, 1913**125 bisulcatus Grandjean, 1953**

Cepheus bifidatus Nicolet, 1855

Damp moss vegetations, Delden, Oudemans; De Lutte, 21.IV.1949, Van der Hammen (1952); Maastricht, St. Pietersberg, 12.IX.1950; Sla- vanste, 25.IX.1951; Plasmolen 22 and 31.X.1968; Westendorp (GE), 22.VI.1980, NAT; Catsop, Elsloo, 2.VI.2000, Zaitsev & Berg (2001); in *Mnium* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993; moss on soil, Mook, Helkuil, 23.VIII.1989, HS.

AMERIDAE**Amerus Berlese, 1896****126 Amerus polonicus Kulczynski, 1902**

Amerus troisi; Sellnick 1928 [misidentification]

Rare species, an old record from Oudemans (Den Haag) in Van der Hammen (1952) and some recent from Zaitsev & Berg (2001).

AMEROBELBIDAE**Amerobelba Berlese, 1908****127 Amerobelba decedens Berlese, 1908**

Eremobelba maxima Willmann, 1951

Adornispolder, Oostburg, 14.VII.1974, Polderman & de Vries (1978); Huys te Linschoten (UT), 4.VI.1985, RMNH; Verlengde Bemerlerberg (LI), 20.XI.2007, HS.

DAMAEOLIDAE**Fosseremus Grandjean, 1954****128 laciniatus (Berlese, 1905)**

Fosseremus quadripertitus Grandjean, 1965

Damaeolus ornatus Mihelcic, 1956

Adornispolder, Oostburg, 14.VII.1974, Polderman & de Vries (1978); Hackfort, Vorden, 1959, Minderman (1981); litter under *Anemone nemorosa*, Maastricht, St. Pietersberg, 21.IV.1994, HS.

CTENOBELBIDAE**Ctenobelba Balogh, 1943****129* pectinigera (Berlese, 1908)**

Murcia obsoleta Koch, 1841 (spec. inquirenda)

Few records: Ooij, Ubbergen (GL), 4.IV.2002; Druten (GL), 6.III.2002, both in grassland on a clay soil; chalk grassland, Eijsderheide, 28.XI.2007, ALT, HS.

EREMAEIDAE**Eueremaeus Mihelcic, 1963****130 oblongus (C.L. Koch, 1835)**

Eremaeus major Mihelcic, 1953

Several records by Van der Hammen (1952) and RMNH.

131* silvestris (Forsslund, 1955)

Eremaeus granulatus Mihelcic, 1955

Eremaeus fuguratus Winkler, 1956

Moss on oak, Hoge Veluwe, Franse Berg, 6.IX.1989, HS; some records under *E. oblongus* (130) may refer to this species.

ZETORCHESTIDAE**Zetorchestes Berlese, 1888****132 flabarius Grandjean, 1951**

Maastricht, St. Pietersberg, 14.X.1952, Krispin (1987).

MICROZETIDAE**Microzetes Berlese, 1913****133 septentrionalis (Kunst, 1963)**

Bovenpolder, Amerongen, 4.IV.2000; Amsterdamse Bos, Amsterdam, 15.VIII.2000, Zaitsev & Berg (2001).

CALEREMAEIDAE

Caleremaeus Berlese, 1910

134 monilipes (Michael, 1882)

Kolhorn remnant in neolithic well, 1983; Schelvis (1992); Winterswijk, 1.VII.1975; Haelen, 28.VI.1978; Platte Bos (LI), 4.VI.1987; Vijlenerbos (LI), 30.III.1975; Bissen (LI), 30.III.1975; Vaals, 28.V.1985; Eperheide (LI); 26.V.1985, RMNH.

TENUIALIDAE

Hafenrefferia Oudemans, 1906

135 gilvipes (C.L. Koch, 1839)

Liacarus pterotus Coggi, 1900
Liacarus auritus Nordenskiöld, 1901
Rare species with few records from Oudemans in Oosterbeek and Arnhem, Van der Hammen (1952); Rhenen, Blauwe Kamer, VI.1991; Doorn, 16.VIII.1983; Winterswijk, 3.VII.1975; Doetinchem, 23.VI.1983; Mook, 8.VI.1988; Valkenburg, 23.V.1982; RMNH; forest litter, Arnhem, Schaarsbergen, 8.XI.1989, HS.

GUSTAVIIDAE

Gustavia Kramer, 1879

136* fusifer (C.L. Koch, 1841)

Winterswijk, 2.VII.1975; Beuningen, 15.VIII.1963; Bemelen, 15.X.1952, RMNH.

137 microcephala (Nicolet, 1855)

Gustavia sol Kramer, 1879
Few records from literature (Leeuwen by Oudemans in Van der Hammen (1952)); recently by Zaitsev & Berg (2001) and RMNH; forest litter, Winterswijk, Bekkendelle, 9.IV.1993, HS.

ASTEGISTIDAE

Astegistes Hull, 1916

138* pilosus (C.L. Koch, 1841)

Acarus muscorum Scopoli, 1763
Zetes pilosus C.L. Koch, 1841
Liacarus bicornis Warburton & Pearce, 1905

Hull (1916) named *Liacarus bicornis* Warburton & Pearce, 1905 the type species of *Astegistes*, which was considered later as a synonym of *Zetes pilosus* C.L. Koch, 1841 and *Acarus muscorum* Scopoli, 1763. Slochteren, 21.VI.1984; Zwartsluis, 13.VI.1985; Groesbeek, 10.VI.1988; Mook, 30.X.1980; Pannerden, 24.VI.1980; Waterland (NH), 5.VI, 3.VII, 6.VII and 19.IX.1984, RMNH; quagmire, Veenendaal, Blauwe Hel, 3.VII.1992, HS.

Cultroribula Berlese, 1908

139 biculturata (Berlese, 1904)

Cultroribula trifurcata Jacot, 1939
Cultroribula szent-ivanovi Balogh, 1943
Cultroribula falcata Evans, 1952
Cultroribula trifurcata rotundata Krivolutski, 1962
Several records by Zaitsev & Berg (2001); forest litter, Utrechtse Bos, 3.V.1994, HS.

140* juncta (Michael, 1885)

Heerenveen, 18.VI.1979; Ubbergen, 7.VI.1988; Trekerduinen (UT), 16.VIII.1983; Onderste Bos (LI), 3.VI.1987; Gulpen, 26.V.1982; Valkenburg, 23.V.1982; Swalmen, 26.VI.1978, RMNH.

LIACARIDAE

Adoristes Hull, 1916

141 ovatus (C.L. Koch, 1839)

Leiosoma pyrigerum Berlese, 1888
Adoristis poppei (Oudemans, 1906)

Several records by Van der Hammen (1952), Zaitsev & Berg (2001), HS, ALT, RMNH
Sub *A. poppei*: Bemelen, 15.X.1952; Platte Bos (LI), 23.VIII.1988, RMNH; oak forest with *Deschampsia flexuosa*, Hoog Soeren, 1.IX.1988; in *Thuja* litter, Beekbergen, Nieuw Dennenlust, 24.VII.1989, HS.

Liacarus Michael, 1898

142 coracinus (C.L. Koch, 1840)

Leiosoma similis Nicolet, 1855
Liacarus parvus Mihelcic, 1956
Several records by Van der Hammen (1952), RMNH.

- 143 nitens (Gervais, 1844)**
Bergh en Terbljijt, Bergheide, 1.vi.2000,
Zaitsev & Berg (2001); Maastricht, St. Pietersberg, 12.IX.1950; Slavante, 25.IX.1951; Roden (DR), 7.IX.1976; Oldematen (ov), 10 and 11.VI.1985, RMNH.
- 144 subterraneus (C.L. Koch, 1844)**
Leiosoma lativentris Nicolet, 1855
Leiosoma globosum Berlese, 1885
Liacarus temellae Willmann, 1931
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH. Hackfort, Vorden, 1959, Minderman (1981); litter between *Equisetum hyemale*, Bekkendelle, Winterswijk, 9.IV.1993, HS.
- 145* xyloiae (Schrank, 1803)**
Groesbeek, 8.VI.1988; Vorden, Hackfort, 1965;
Heerenveen, 18.VI.1979; Valkenburg, 23.V.1982, RMNH; old hedgerow, Harskamp, 25.IX.1997, HS.
- Xenillus Robineau-Desvoidy, 1839**
- 146 clypeator Robineau-Desvoidy, 1839**
Xenillus latus (Nicolet, 1855)
Cepheus permixtus André, 1925
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, HS.
- 147* discrepans Grandjean, 1936**
Moss on tree trunk, Mook, De Diepen, 23.VIII.1989, HS.
- 148 tegeocranus (Hermann, 1804)**
Cepheus vulgaris Nicolet, 1855
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH. Hackfort, Vorden, 1959, Minderman (1981); forest litter, Arnhem, Schaarsbergen, 20.VII.1989, HS.
- CARABODIDAE**
- Carabodes C.L. Koch, 1835**
- 150* areolatus Berlese, 1916**
Westerbroek (GR), 20.VI.1984; Slochteren, 21.VI.1984; Winterswijk, 1.VII.1975; Apeldoorn, V.1984; Vredepeel (NB), 31.OCT.1975; Valkenburg, 23.V.1982; Vijlenerbos (LI), 29.V.1974, 30.III.1975; Bissen (LI), 30.III.1975, RMNH.
- 151 coriaceus C.L. Koch, 1835**
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; hedgerow, Klarenbeek, 22.IX.1997, HS.
- 152 femoralis (Nicolet, 1855)**
Several records by Zaitsev & Berg (2001) and RMNH; forest litter, Ugchelense Bos, 3.V.1994, HS.
- 153 labyrinthicus (Michael, 1879)**
Cepheus heimi Oudemans, 1903
Carabodes vermiculatus Berlese, 1916
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.
- 154 marginatus (Michael, 1890)**
Cepheus minutus Oudemans, 1896
Several records by Van der Hammen (1952) and RMNH; grassy heathland, Deelerwoud, 22.VIII.1988, HS.
- 155* ornatus Storkan, 1925**
Carabodes forsslundi Sellnick, 1953
Bissen (LI), 30.III.1975, Winterswijk, 1.VII.1975, RMNH.
- 156* rugosior Berlese, 1916**
Vijlenerbos (LI), 29.V.1974, RMNH.
- 157 willmanni Bernini, 1975**
Carabodes minusculus; Sellnick 1928,
Willmann 1931 [misidentification]
Several records by Van der Hammen (1952) and Zaitsev & Berg (2001); in *Cladonia arbuscula* in sand dunes, Texel, De Slufter, 7.II.1993, HS. Many of the older records of *C. minusculus* refer to this species; *C. willmanni* is recognised since 1975.
- METRIOPPIIDAE**
- Ceratoppia Berlese, 1908**
- 149 bipilis (Hermann, 1804)**
Ceratoppia herculeana Berlese, 1908
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.

Odontocepheus Berlese, 1913
158 elongatus (Michael, 1879)

Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; forest litter,
Hoenderloose Bos, 3.v.1994, HS.

TECTOCEPHEIDAE

Tectocepheus Berlese, 1896

159* alatus Berlese, 1913

Texel, De Mok, 14.XII.1974, RMNH.

160* bisignatus Moritz, 1968

In *Marchantia* at a steep slope of a stream,
Döttekro, Winterswijk, 9.IV.1993, HS.

161 cuspidentatus Knülle, 1954

Several records by Zaitsev & Berg (2001).

162* sarekensis Trägårdh, 1910

Tectocepheus velatus sarekensis Trägårdh,
1910

Tectocepheus velatus angulatus Mihelcic,
1957

Tectocepheus velatus ibericus Mihelcic, 1957

Tectocepheus velatus inflexus Mihelcic, 1957
In *Mnium* on border of a ditch, Beekbergen,
Elsbosweg, 14.VII.1991, HS.

163 velatus (Michael, 1880)

Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.

QUADROPPIIDAE

Quadroppia Jacot, 1939

164 bellula Luxton, 1987

Several records by Zaitsev & Berg (2001).

165* maritalis Lions, 1982

Moss under *Sambucus*, Texel, Horsmeertjes,
6.II.1993, HS.

166 michaeli Mahunka, 1977

Several records by Zaitsev & Berg (2001).

167* monstruosa Hammer, 1979

Among litter with *Anemone nemorosa*,
St. Pietersberg, Maastricht, 21.IV.1994, HS.

168 quadricarinata (Michael, 1885)

Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; soil sample
(depth: 4.6 cm), Deelerwoud, 24.VI.1988, HS.
Note: Some of the older records may refer to
the more recently described *Quadroppia* sp.

MACHUELLIDAE

Machuella Hammer, 1961

169 draconis Hammer, 1961

Several records by Zaitsev & Berg (2001);
birchbrook, Zwilbroek, Eibergen, 9.IV.1993,
HS.

OPPIIDAE

Berniniella Balogh, 1983

170* bicarinata (Paoli, 1908)

Nijmegen, v.1976, moss in garden, RMNH.

171* lunaris (Evans, 1952)

Litter between *Equisetum hyemale*,
Bekkendelle, Winterswijk, 9.IV.1993, HS.

172 serratirostris (Golosova, 1970)

Zwilbroek, Eibergen, 16.III.1999, Zaitsev &
Berg (2001).

173 sigma (Strenzke, 1951)

Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001).

Dissorrhina Hull, 1916

174 ornata (Oudemans, 1900)

Damaeosoma tricarinata Paoli, 1908

Damaeosoma splendens; Berlese 1895 [mis-
identification]

Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.

Hypogeoppia Subias, 1981

175* terricola Subias, 1981

Old hedgerow, Epe, Vemde, 22.IX.1997, HS.

Ramusella Hammer, 1962

176 clavipectinata (Michael, 1885)

Several records by Van der Hammen (1952),
ALT and RMNH.

- 177* *furcata* (Willmann, 1928)
Oppia furcata; Kunst 1958 [misidentification]
Linschoten (UT), 4.VI.1985, RMNH.
- 178* *insculpta* (Paoli, 1908)
Old hedgerow, Woudenberg, 25.IX.1997, hs.
- Micropippia Balogh, 1983**
- 179 *minus* (Paoli, 1908)
Several records by Zaitsev & Berg (2001) and RMNH; grassland, Hoge Veluwe, Compagnieberg 1.VIII.1989, hs.
- 180* *minutissima* (Sellnick, 1950)
Forest litter, Ugchelense Bos, 3.V.1994, hs.
- Moritzoppia Subias & Minguez, 1988**
- 181 *neerlandica* (Oudemans, 1900)
Eremaeus longilamellatus var. *neerlandica* Oudemans, 1900
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH.
- 182* *translamellata* (Willmann, 1923)
Moss on tree trunk, Leersum, VI.1975, RMNH.
- 183 *unicarinata* (Paoli, 1908)
Mantingerbos, Mantinge, 23.VII.1949, Van der Hammen (1952); in moss at the foot of an oak, Rhenen, Grebbeberg, IX.1989, hs; Rhenen, Blauwe Kamer, VI.1991; Weert, 1976; Westerbroek (GR), 20.VI.1984; Slochteren, 21.VI.1984, RMNH.
- Oppia C.L. Koch, 1835**
- 184 *nitens* C.L. Koch, 1835
Maastricht, St. Pietersberg, 6.V.1949, Van der Hammen (1952).
- Oppiella Jacot, 1937**
- 185 *falcata* (Paoli, 1908)
Several records by Van der Hammen (1952) and RMNH; old hedgerow, Epe, Vemde, 22.IX.1997, hs.
- 186* *maritima* (Willmann, 1929)
Damaeosoma falcatum maritima Willmann, 1929
- Oppia fissurata*, Hammer, 1952
In moss, Winterswijk, Willink Weuste, 9.IV.1993, hs.
- 187 *nova* (Oudemans, 1902)
Damaeosoma corrugatum Berlese, 1904
Notaspis sculptilis Warburton & Pearce, 1905
Damaeosoma neerlandicum; Sellnick 1928 [misidentification]
- Oppia neerlandica*; Willmann 1931 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001), hs, ALT and RMNH.
- 188* *splendens* (C.L. Koch, 1841)
Vaals, 28.V.1982; Wrakelberg (LI), 23.VIII.1988; Vijlenerbos (LI), 22.VIII.1988; Platte Bos (LI), VI.1986; Eyserbos (LI), VI.1986; Gulpen, 26.V.1982, RMNH; beech forest east of Nijmegen, 3.IX.1989, hs.
- 189* *uliginosa* (Willmann, 1919)
Birchbrook, Zwilbroek, Eibergen, 9.IV.1993, hs.
- Rhinoppia Balogh, 1983**
- 190* *fallax* (Paoli, 1908)
Oppiella dubia Hammer, 1962
Heerenveen, 18.VI.1979; Gaasterland, 19.VI.1979; Waterland (NH), 12.VI.1984; Eyserbos (LI), 2.VI.1987; St.Odiliënberg, 29.VI.1978; Vaals, 28.V.1982; Gulpen, 26.V.1982, RMNH.
- 191 *obsoleta* (Paoli, 1908)
Damaeosoma vitrinum Hull, 1914
Several records by Zaitsev & Berg (2001) (sub *M. media*) and RMNH; in *Rhytidiaadelphus squarrosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, hs.
- 192 *subpectinata* (Oudemans, 1900)
Belba pectinata; Berlese 1892 [misidentification]
- Damaeosoma clavipectinatum*; Paoli 1908 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Mnium* at a steep slope of a stream, Döttekörö, Winterswijk, 9.IV.1993, hs.

SUCTOBELBIDAE

- Allosuctobelba Moritz, 1970**
193* *grandis* (Paoli, 1908)
Suctobelba grandis europaea Willmann, 1933
Suctobelba europaea Willmann, 1933
Zeevang, 21.vi.1983; Schoorl, 21.vi.1983;
Gulpen, 26.v.1982, RMNH.
- Suctobelba Paoli, 1908**
194* *altvateri* Moritz, 1970
In *Mnium* on border of a ditch, Beekbergen,
Elsbosweg, 14.VII.1991, HS.
- 195 *granulata* Van der Hammen, 1952
Suctobelba trigona f. *granulata* Van der
Hammen, 1952
Several records by Van der Hammen (1952);
in *Mnium* on border of a ditch, Beekbergen,
Elsbosweg, 14.VII.1991, HS.
- 196* *secta* Moritz, 1970
Litter between *Equisetum hyemale*,
Bekkendelle, Winterswijk, 9.IV.1993, HS.
- 197* *trigona* (Michael, 1888)
Hasselt, 11.VI.1985; Ubbergen, 7.VI.1988;
Trekkerduinen (UT), 16.VIII.1983; Spaubeek,
25.V.1982; Valkenburg, 23.V.1982; Mook,
8.VI.1988; Onderste Bos (LI), 3.VI.1987;
Wellerlooise Bos (LI), 1977, RMNH.
- Suctobelbellula Jacot, 1937**
- 198 *acutidens* (Forsslund, 1941)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; forest litter,
Hoenderloose Bos, 3.V.1994, HS.
- 199 *lobata* (Strenzke, 1950)
Suctobelba acutidens lobata Strenzke, 1950
Suctobelbellula acutidens lobata (Strenzke, 1950)
Several records by Zaitsev & Berg (2001); pine
litter, Ysselstein (LB), 28.III.1993, HS.
- 200 *alloenasuta* Moritz, 1971
Several records by Zaitsev & Berg (2001); old
hedgerow, Klarenbeek, 22.IX.1997, HS.
- 201 *falcata* (Forsslund, 1941)
Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001); forest litter,
Ugchelense Bos, 3.V.1994, HS.
- 202 *forsslundi* (Strenzke, 1950)
Several records by Zaitsev & Berg (2001);
under *Allium ursinum*, Rijkholt, Savelbos,
21.IV.1994, HS.
- 203* *hamata* Moritz, 1970
In *Marchantia* at a steep slope of a stream,
Döttekro, Winterswijk, 9.IV.1993, HS.
- 204 *hammerae* (Krivolutski, 1965)
Suctobelba hammeri Krivolutski, 1965
Suctobelba sarekensis; Hammer 1952 [mis-
identification]
Several records by Zaitsev & Berg (2001).
- 205 *longirostris* (Forsslund, 1941)
Several records by Zaitsev & Berg (2001).
- 206* *nasalis* (Forsslund, 1941)
In moss, Winterswijk, Willink Weuste,
9.IV.1993, HS.
- 207 *palustris* (Forsslund, 1953)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; quagmire,
Veenendaal, Blauwe Hel, 3.VII.1992, HS.
- 208 *perforata* (Strenzke, 1950)
Several records by Zaitsev & Berg (2001) and
RMNH.
- 209 *sarekensis* (Forsslund, 1941)
Suctobelba cornigera; Trägårdh 1910 [mis-
identification]
Several records by Van der Hammen (1952)
and Zaitsev & Berg (2001).
- 210 *similis* (Forsslund, 1941)
Several records by Zaitsev & Berg (2001); for-
est litter, Ugchelense Bos, 3.V.1994, HS.
- 211 *subcornigera* (Forsslund, 1941)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS and ALT.
- 212 *subtrigona* (Oudemans, 1900)
Suctobelba intermedia Willmann, 1939
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; pine litter,
Ysselstein (LI), 28.III.1993, HS.
- 213* *tuberculata* (Strenzke, 1950)
Boswachterij Ruurlo (GE), 14.IX.2007; Groll-
oërveld (GE), 21.IX.2007, HS.
- 214 *vera* (Moritz, 1964)
Several records by Zaitsev & Berg (2001);
forest litter, Ugchelense Bos, 3.V.1994, HS.

AUTOGENETIDAE**Autogneta Hull, 1916**

- 215
- longilamellata*
- (Michael, 1888)

Several records by Van der Hammen (1952) and Zaitsev & Berg (2001); Hackfort, Vorden, 1959, Minderman (1981); Ameland, Nesserbos, 28.VI.1972, RMNH; birchbrook, Zwilbroek, Eibergen, 9.IV.1993, HS.

Conchogneta Grandjean, 1963

- 216
- delacarlica*
- (Forsslund, 1947)

Autogneta willmanni (Dyrdowska, 1929)
Several records by Van der Hammen (1952) and Zaitsev & Berg (2001); litter under *Ficaria verna*, Winterswijk, Bekkendelle, 9.IV.1993, HS.

THYRISOMIDAE**Banksinoma Oudemans 1930**

- 217
- lanceolata*
- (Michael, 1885)

Oribella castanea; Willmann 1931 [misidentification]
Xenillus castaneus; Sellnick 1928 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Rhytidia-delphus squarosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, HS.

- 218
- oudemansi*
- Fujikawa, 1978

A record from Oudemans, Haarlem, VIII.1898, Fujikawa (1978).

Oribella Berlese, 1908

- 219
- pectinata*
- (Michael, 1885)

Several records by Van der Hammen (1952); *Sphagnum* in forest, Groesbeek, 7.XI.1989, HS.

Pantelozetes Grandjean, 1953

- 220*
- multidentata*
- (Evans, 1954)

Oribella paolii multidentata Evans, 1954
In moss, Winterswijk, Willink Weuste, 9.IV.1993, HS.

- 221
- paolii*
- (Oudemans, 1913)

Eremaeus pectinatus; Oudemans 1900 [misidentification]

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, HS.

HYDROZETIDAE**Hydrozetes Berlese, 1902**

- 222
- confervae*
- (Schrank, 1781)

Hydrozetes terrestris Berlese, 1910
Cepheus (Carabodes) coriaceus; Oudemans 1896 [misidentification]
Records from Oudemans at Sneek and Utrecht, Van der Hammen (1952); quagmire, Westbroekse Zodden, 29.VI.1992, HS.

- 223
- lacustris*
- (Michael, 1882)

Notaspis speciosus Chinaglia, 1917
Several records by Van der Hammen (1952); Losser, 1 and 3.VI.1990, Waterland (NH), 3.VII.1984, RMNH; *Sphagnum*, Hoge Veluwe, IJzeren Man, 16.VIII.1990, HS.

- 224
- lemnae*
- (Coggi, 1899)

A record from Valkveen found in the Oudemans collection by Van der Hammen (1952); Kamperland, Veerse Meer, 5.IX.1975, RMNH.

- 225*
- parisiensis*
- Grandjean, 1948

Maarssen, Molenpolder, Westbroek, 3.V.1972, RMNH.

LIMNOZETIDAE**Limnozetes Hull, 1916**

- 226
- ciliatus*
- (Schrank, 1803)

Oribata sphagni Michael, 1884
Several records by Van der Hammen (1952) and RMNH; quagmire, Westbroekse Zodden 29.VI.1992, HS.

227* foveolatus Willmann, 1939

Limnozetes ciliatus foveolatus Willmann, 1939
Limnozetes palmerae Behan-Pelletier, 1989
Limnozetes sphagni; Sellnick 1928 [misidentification]
In pond with *Sphagnum*, Beekbergen, Reeënberg, 19.VII.1989, hs.
Due to the former confusion with *L. ciliatus*, many records of *L. foveolatus* may be listed under this species.

228 rugosus (Sellnick, 1923)

Several records by Van der Hammen (1952) and RMNH; *Sphagnum*, Hoge Veluwe, IJzeren Man, 16.VIII.1990, hs.

AMERONOTHRIDAE

Ameronothrus Berlese, 1896

229 maculatus (Michael, 1882)

Scutovertex pseudomaculatus Hull, 1914
Records from the sea shore at Zeeburg and inland in Oude Pekela from Oudemans (Van der Hammen (1952) and from the bark of *Platanus* (Noordijk & Berg 2001); many records from the seashore in RMNH.

230 marinus (Banks, 1896)

Scutovertex spoofi Oudemans, 1900
Scutovertex bilineatus; Oudemans 1901 [misidentification]
Ameronothrus bilineatus spoofi Hull, 1916
Ameronothrus fucicolum; Buitendijk 1945 [misidentification]
Ameronothrus infusorium; Buitendijk 1945 [misidentification]
A record from Oudemans at Zeeburg at the sea shore, Van der Hammen (1952); basalt dike, Dollard, Uithuizerwad, 1.XI.1974, RMNH.

231* nigrofemoratus (L. Koch, 1879)

Ameronothrus lineatus var. nigrofemoratus (L. Koch, 1879)
Ameronothrus lineatus brevipes Willmann, 1937
Many records from the seashore in RMNH.

232 schneideri (Oudemans, 1905)

Many records from the seashore in RMNH (Polderman 1974).

CYMBAEREMAEIDAE

Cymbaeremaeus Berlese, 1896

233 cymba (Nicolet, 1855)

Cymbaeremaeus pulvillifer Willmann, 1931 (nymph)
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; litter around *Juniperus* shrubs, Otterlo, 7.VIII.1989, hs.

Scapheremaeus Berlese, 1910

234 palustris Sellnick, 1924

One old record from Oudemans collected from a pond (Van der Hammen (1952)).

MICREREMIDAE

Micreremus Berlese, 1896

235 brevipes (Michael, 1888)

Several records by Van der Hammen (1952) and RMNH; litter between *Equisetum hyemale*, Bekkendelle, Winterswijk, 9.IV.1993, hs.

LICNEREMAEIDAE

Licneremaeus Paoli, 1908

236 licnophorus (Michael, 1882)

Arnhem, 9.IX.1965; Vriezenveen, 1974; Hilvarenbeek, 6.VI.1977, RMNH; Litter around *Juniperus* shrubs, Otterlo, 7.VIII.1989, hs.

PASSALOZETIDAE

Passalozetes Grandjean, 1932

237 strenzkei Weigmann, 2006

Passalozetes bidactylus; Willmann 1931, Strenzke 1953 [misidentification]
Records from Oudemans at Valkeveen and at Meyendel, 24.III.1949, Van der Hammen (1952); Domburg, v.1977, Texel, Mokbaai, 12.VIII.1976; Ameland, Buren, 8.VI.1983; Siebengewald (LI), 1977, RMNH; in *Rhytidiaadelphus squarrosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, hs.

- 238* **perforatus** (Berlese, 1910)
Wijk aan Zee (NH), de Rellen and Vogelduinen, 24.X.2007, HS.
- SCUTOVERTICIDAE**
- Scutovertex** Michael, 1879
- 239 **minutus** (C.L. Koch, 1836)
Scutovertex ovalis Oudemans, 1900
Several records by Van der Hammen (1952),
RMNH, Arnhem Schaarsbergen Forest litter
14.VII.1989, HS.
- 240 **pilososetosus** Polderman, 1977
Polderman (1977).
- 241 **sculptus** Michael, 1879
Several records in Van der Hammen (1952)
and RMNH.
- PELOPSIDAE**
- Eupelops** Ewing, 1917
- 242 **acromios** (Hermann, 1804)
Pelops planicornis Willmann, 1931
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH (sub *P. planicornis*); forest litter, Arnhem, Schaarsbergen,
31.VIII.1989, HS.
- 243 **curtipilus** (Berlese, 1916)
Eupelops bilobus (Sellnick, 1928)
Cladonia in heathland, Hoge Veluwe,
Hoenderloo, 11.X.1948; moss in heathland,
Ossendrecht, 23.II.1949, Van der Hammen
(1952).
- 244 **occultus** (C.L. Koch, 1836)
Eupelops nepotulus; Sellnick 1960 [misidentification]
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; in *Rhytidadelphus squarosus* on sand dunes, Horsmeertjes,
Texel, 6.II.1993; old hedgerow, Klarenbeek,
22.IX.1997, HS.
- 245 **oudemansi** (Van der Hammen, 1952)
Record from the type locality Valkeveen in the
Oudemans collection (Van der Hammen
1952).
- 246 **plicatus** (C.L. Koch, 1835)
Pelops auritus C.L. Koch, 1839
Pelops laevigatus Nicolet, 1855
Pelops fuligineus; Oudemans 1896 [misidentification]
Pelops fusiger Mihelcic, 1957
Pelops acromios diversipilus Mihelcic, 1957
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH (part. sub
P. auritus); Hackfort, Vorden, 1959, Minder-
man (1981); litter under *Ficaria verna*, Winters-
wijk, Bekkendelle, 9.IV.1993, HS.
- 247* **strenzkei** (Knüllé, 1954)
Quagmire, Westbroekse Zodden, 29.VI.1992,
HS.
- 248* **tardus** (C.L. Koch, 1836)
Moss under *Sambucus*, Texel, Horsmeertjes,
6.II.1993, HS.
- 249 **torulosus** (C.L. Koch, 1836)
Rhedense Heide, Rheden, 28.VIII.2000,
Zaitsev & Berg (2001).
- 250 **peloptulus** Berlese, 1908
- 250* **montanus** Hull, 1914
Probably subfossil records only (peat core
Vriezenveen), RMNH.
- 251 **phaenotus** (C.L. Koch, 1844)
Several records by Van der Hammen (1952)
and RMNH; in *Rhytidadelphus squarosus* on
sand dunes, Horsmeertjes, Texel, 6.II.1993, HS.
- ACHIPTERIIDAE**
- Achipteria** Berlese, 1885
- 252 **coleoptrata** (Linnaeus, 1758)
Oribata ovalis Nicolet, 1855
Oribates nicoletii Berlese, 1883
Oribata intermedia Michael, 1898
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; Hackfort,
Vorden, 1959, Minderman (1981); in *Rhytidadelphus squarosus* on sand dunes, Horsmeertjes,
Texel, 6.II.1993, HS.

- 253 nitens (Nicolet, 1855)**
Notaspis mixtus Sellnick, 1943
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981), in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, hs.
- Parachipteria Van der Hammen, 1952**
- 254 punctata (Nicolet, 1855)**
Notaspis italicus; Willmann 1931 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; litter around *Juniperus* shrubs, Otterlo 6.IX.1989, hs.
- 255 willmanni Van der Hammen, 1952**
Notaspis punctatus; Willmann 1931 [misidentification]
Several records by Van der Hammen (1952) and RMNH; in *Mnium undulatum* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, hs.
- Pseudachipteria Travé, 1960**
- 256* magna (Sellnick, 1928)**
Hilversumse Wasmeer, 22.VI.2000, hs.
- ORIBATELLIDAE**
- Ophidiotrichus Grandjean, 1953**
- 257 tectus (Michael, 1884)**
Notaspis borussicus Sellnick, 1909
Tectoribates connexus var. borussicus Sellnick, 1928 (sec. homonym)
Joelia connexa var. borussica (Sellnick, 1928) (sec. homonym)
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH (sub *J. borussica*); Hackfort, Vorden, 1959, Minderman (1981); forest litter, Ugchelense Bos, 3.V.1994, hs.
- Oribatella Banks, 1895**
- 258* arctica littoralis Strenzke, 1950**
Several records from the seashore in RMNH; in grass litter, Texel, De Slufter, 7.II.1993, hs.
- 259 berlesei Michael, 1898**
Several records by Zaitsev & Berg (2001).
- 260 calcarata (C.L. Koch, 1835)**
Oribatella decumana Berlese, 1910
Several records by Zaitsev & Berg (2001) and RMNH; moss on tree trunk, Ittervoort, Vijverbroek, 28.III.1993, hs.
- 261 quadricornuta (Michael, 1880)**
Oribates flammula; Oudemans 1896 [misidentification]
Oribatella calcarata; Willmann 1931 [misidentification]
Several records by Van der Hammen (1952) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); litter of *Thuja*, Beekbergen, Nieuw Dennenlust, 11.VII.1989, hs.
- 262* reticulata Berlese, 1916**
Oak forest, east of Nijmegen, 24.IX.1989, hs.
- 263* superbula (Berlese, 1904)**
Oribatella meridionalis Berlese, 1908
Oribatella willmanni (Subias & Gil-Martin, 1995)
Heerenveen, 18.VI.1979; Holwerd (FR), 20.VI.1979; Doetinchem, 23.VI.1980; Waterland (NH), 3.VII.1984; St. Odiliënberg (LI), 29.VI.1978, RMNH (sub *O. meridionalis*).
- GALUMNIDAE**
- Acrogalumna Grandjean, 1956**
- 264 longipluma (Berlese, 1904)**
Oribates elimatus longiplanus Berlese, 1904
Oribata setiformes Hall, 1911
Galumna filata Oudemans, 1913
Oribates elimatus; Oudemans 1896 [misidentification]
Notaspis elimatus; Oudemans 1900 [misidentification]
Several records by Van der Hammen (1952) and Zaitsev & Berg (2001); Huys te Linschoten (UT), 4.VI.1985 and more sub *G. longipluma*, RMNH; in *Mnium undulatum* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, hs.

Allogalumna Grandjean, 1936**265 neerlandica Van der Hammen, 1952**

Texel (type locality), 7.v.1949, Van der Hammen (1952); Texel, 3.iv.1974, 1.i.1971 and 11.vi.1974; Meyendel (ZH), 21.i.1974; Slochteren, 21.vi.1984, RMNH; in *Rhytidia delphus squarosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, hs.

Galumna von Heyden, 1826**266 alata (Hermann, 1804)**

Rockanje, 17.IX.1948, Van der Hammen (1952); Boetwetering, Windesheim, 6.II.2000, (Zaitsev & Berg 2001); Texel, Polder Eendracht, 1.VII.1971; Texel, Polder het Noorden, 13.IV.1972, RMNH; moss under *Sambucus*, Texel, Horsmeertjes, 6.II.1993, hs.

267 lanceata (Oudemans, 1900)

Notaspis alatus; Oudemans 1901 [misidentification]
Galumna dorsalis; Van der Drift 1950 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; *Cirsio-Molinietum*, Zegveld, 22.VI.1992, hs.

268 obvia (Berlese, 1913)

Galumna elimata; Willmann 1931 [misidentification]
Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; among litter, Texel, Mokbaai, 6.II.1993, hs.

Pergalumna Grandjean, 1936**269 nervosa (Berlese, 1914)**

Galumna retalata Oudemans, 1915
Zetes nervosus (Berlese, 1914)
Several records by Van der Hammen (1952) and RMNH; in moss, Winterswijk, Willink Weuste, 9.IV.1993, hs.

270* willmanni (Zachvatkin, 1953)

Galumna formicaria; Willmann 1931 [misidentification]
Doetinchem, 23.VI.1980; Ravensbos (LI), 23.V.1982; Waterop (LI), 26.V.1982, RMNH (sub *G. formicarius*).

Pilogalumna Grandjean, 1956**271* crassiclava (Berlese, 1914)**

Galumna allifera Oudemans, 1915
Wieringen (NH), 30.VIII.1983, Wrakelberg (LI), 23.VIII.1988, RMNH (sub *G. allifera*).

272 tenuiclava (Berlese, 1908)

Galumna areolata Willmann, 1923
Galumna radiata Sellnick, 1928

Several records by Van der Hammen (1952) and RMNH; litter of *Thuja*, Beekbergen, Nieuw Dennenlust, 17.VII.1989, hs.

CERATOZETIDAE**Ceratozetes Berlese, 1908****273 gracilis (Michael, 1884)**

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, hs.

274 mediocris Berlese, 1908

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH.

275* parvulus Sellnick, 1922

Several records in RMNH but probably all subfossil (from peat cores).

276* peritus Grandjean, 1951

Adornispolder (ZE), 14.VII.1974, RMNH.

Diapterobates Grandjean, 1936**277 humeralis (Hermann, 1804)**

Sphaerozetes (Trichoribates) numerosus Sellnick, 1924

Oribates picipes Hull, 1913

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); Veenendaal, Blauwe Hel, 31.VIII.1989, hs.

- Edwardzetes Berlese, 1914**
- 278 edwardsi (Nicolet, 1855)**
- Oribata edwardsi Nicolet, 1855
 - Maastricht, St. Pietersberg, 7.v.1949, Van der Hammen (1952); Winterswijk, Korenburgrerveen, 2.vii.1975; Ubbergen, 27.vi.1980; Nijmegen, 20.iv.1974; Gerendal (LI), 5.v.1972; Ravensbos (LI), 27.v.1974, RMNH; in moss, Beuningen, Het Broek, 31.viii.1989, HS.
- Fuscozetes Sellnick, 1928**
- 279 fuscipes (C.L. Koch, 1844)**
- Galumna slossonae Banks, 1906
 - Fuscozetes rotundatus Willmann, 1930
 - Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Mnium undulatum* at a steep slope of a stream, Döttekro, Winterswijk, 9.iv.1993, HS.
- Latilamellobates Shaldubina, 1971**
- 280 incisellus (Kramer, 1897)**
- Several records by Van der Hammen (1952) and RMNH; in *Rhytidiaadelphus squarosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, HS.
- Melanozetes Hull, 1916**
- 281 mollicomus (C.L. Koch, 1839)**
- Several records by Van der Hammen (1952) and RMNH; moss on soil, Mook, Helkuil, 23.VIII.1989, HS.
- Trichoribates Berlese, 1910**
- 282 novus (Sellnick, 1928)**
- Oribata setosa; Michael 1884 [misidentification]
 - Trichoribates numerosa; Buitendijk 1945 [misidentification]
 - Several records by Van der Hammen (1952) and RMNH; lawn, Beekbergen, Nieuw Denne lust, 24.X.1989, HS.
- 283 trimaculatus (C.L. Koch, 1835)**
- Oribates setosus; Berlese 1887 [misidentification]
 - Sphaerozetes (Trichoribates) berlesei Jacot, 1929
 - Several records by Van der Hammen (1952) and RMNH; *Lolium*-grassland, Arnhem, Bakenbergseweg, 14.XI.1988, HS.
- HUMEROBATIDAE**
- Humerobates Sellnick, 1928**
- 284 rostrolamellatus Grandjean, 1936**
- Oribates humeralis; Berlese 1883 [misidentification]
 - Oribata lapidaria; Michael 1884 [misidentification]
 - Oribates facula; Oudemans 1896 [misidentification]
 - Notaspis humeralis; Oudemans 1902 [misidentification]
 - Humerobates humeralis; Sellnick 1928 [misidentification]
 - Humerobates fungorum; Willmann 1931 [misidentification]
 - Banksinus arboreus rostrolamellatus Jacot, 1940
 - Several records by Van der Hammen (1952), Zaitsev & Berg (2001), RMNH and Noordijk & Berg (2001); on *Calluna vulgaris*, Terlet, 11.IX.1989, HS.
- ZETOMIMIDAE**
- Heterozetes Willmann, 1917**
- 285 palustris Willmann, 1917**
- Doetinchem, Oudemans collection, Van der Hammen (1952); Hoge en Lage Mierde, De Utrecht (NB), 15.VI.1977; Groesbeek, De Bruuk, 10.VI.1988, RMNH; quagmire, Westbroekse Zodden, 29.VI.1992, HS.
- Zetomimus Hull, 1916**
- 286* furcatus (Pearce & Warburton, 1905)**
- Quagmire, Westbroekse Zodden, 29.VI.1992, HS.

CHAMOBATIDAE**Chamobates Hull, 1916**287 *borealis* (Trägårdh, 1902)Notaspis cuspidata var. *borealis* Trägårdh, 1902

Chamobates incisus Van der Hammen, 1952

Chamobates schuetzi; Sellnick 1929 [misidentification]

Chamobates schuetzi; Willmann 1931 [misidentification]

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Thuja* litter, Beekbergen, Nieuw Dennenlust, 24.VII.1989, HS.288 *cuspidatus* (Michael, 1884)

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; birchbrook, Zwilbroek, Eibergen, 9.IV.1993, HS.

289* *pusillus* (Berlese, 1895)

Heerenveen, 18.VI.1979; Opsterland, 21.VI.1979; Denekamp, 30.V.1975; Vriezenveen, 1974; Treekerduinen (UT), 16.VIII.1983; Hoge en Lage Mierde, De Utrecht (NB), 15.VI.1977; Hilvarenbeek, 16.VI.1977; Epen, Bovenste Bos (LI), 16.IX.1951, RMNH; moss on soil, Mook, Helkuil, 23.VIII.1989, HS.

290 *schuetzi* (Oudemans, 1902)Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; on *Calluna vulgaris*, Terlet, 11.IX.1989, HS.291 *subglobulus* (Oudemans, 1900)

Oribata lapidaria Lucas, 1849

Oribata omissa Warburton & Pearce, 1905

Several records by Van der Hammen (1952) and RMNH (partly sub *C. lapidarius*); between *Mnium undulatum*, Haelen, Leudal, 8.X.1989, HS.**Globozetes Sellnick, 1928**292* *longipilus* Sellnick, 1928

Winterswijk, Buskersbos, 2.VII.1975, RMNH; Deelerwoud, soil sample (depth 4.6 cm), 14.I.1988, HS.

293* *tricuspidatus* (Willmann, 1953)In *Mnium undulatum* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, HS.**MYCOBATIDAE****Minunthozetes Hull, 1916**294* *pseudofusiger* (Schweizer, 1922)

Puncitoribates atomus Berlese, 1908 (spec. inquirendae)

Groesbeek, 8 and 10.VI.1988; Hoge en Lage Mierde, De Utrecht (NB), 15.VI.1977; Wrakelberg (LI), VI.1986; Platte Bos (LI), VI.1986, RMNH.

295 *semirufus* (C.L. Koch, 1841)

Oribata fusigera Michael, 1884

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, HS.**Mycobates Hull, 1916**296 *parmeliae* (Michael, 1884)

An old record from Nijkerk in the Oudemans collection (Van der Hammen 1952).

Puncitoribates Berlese, 1908297* *hexagonus* Berlese, 1908

Ameland, Westergie, 19.X.1975; Texel, De Bol, 13.IV.1971; Wervershoof (NH), 30.VIII.1983; Broek in Waterland, 5.VI.1984; Goeree, Springerborg, 21.IX.1973 and 15.VI.1974; Zuid-Beveland, Rilland, 28.VIII.1975; St. Philipsland, 28.VIII.1975; Tholen, Stavenisse, 2.XI.1974; Vleuten, VIII.1976; Nijmegen, Lankhorst, 20.V.1975; St. Odiliënberg (LI), 29.VI.1978; Melick, Herkenbosch, 27.VI.1978, RMNH.

298 *punctum* (C.L. Koch, 1839)

Oribata avenifera Michael, 1884

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); lawn, Beekbergen, Nieuw Dennenlust, 24.X.1989, HS.

299 sellnicki Willmann, 1928

Punc toribates bicornis Sellnick, 1928

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in pond with *Sphagnum*, Beekbergen, Reeënberg, 19.VII.1989, hs.

Zachvatkinibates Shaldybina, 1973**300* quadrivertex (Halbert, 1920)**

Several records in RMNH from the seashore; in grass litter, Texel, De Slufter, 7.II.1993, hs.

EUZETIDAE**Euzetes Berlese, 1908****301 globulus (Nicolet, 1855)**

Carabodes nitens Johnston, 1853

Oribates seminulum Oudemans, 1896

Euzetes aterrimus Sellnick, 1928

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; Hackfort, Vorden, 1959, Minderman (1981); in *Mnium undulatum* at a steep slope of a stream, Döttekro, Winterswijk, 9.IV.1993, hs.

HAPLOZETIDAE**Protoribates Berlese, 1908****302* capucinus Berlese, 1908**

Oribata oblonga Ewing, 1909

Maastricht, St. Pietersberg, 7.V.1949, Van der Hammen (1952); Oldematen (ov), 10.VI.1985, RMNH; Boerwetering, Windesheim, 6.II.2000, Zaitsev & Berg (2001); Colmont, Wrakelberg (LI), 23.XI.2008; Eijserheide (LI), 28.XI.2008; Bemelen, Bemerlerberg, 28.XI.2008, ALT.

303* lophotrichus (Berlese, 1904)

Oldematen (ov), 11.VI.1985; Wrakelberg (LI), 23.VIII.1988, RMNH; grassland on peat soil, Zegveld (UT), 19.IV.2004, ALT.

SCHELORIBATIDAE**Dometorina Grandjean, 1951****304 plantivaga (Berlese, 1895)**

Oppia tibialis; Berlese 1892 [misidentification]

An old record from Arnhem in the Oudemans collection, Neercanne, 7.V.1949, Van der Hammen (1952); Starnuman Bossen, Gaasterland, 24.IV.2000, Zaitsev & Berg (2001); Winterswijk, 1.VII.1975; Putberg (LI), VI.1986, RMNH; between *Mnium undulatum*, Haelen, Leudal, 8.X.1989, hs.

Hemileius Berlese, 1908**305 initialis (Berlese, 1908)**

Scheloribates confundatus Sellnick, 1928

Hemileius translamellatus; Karpen 1974 [misidentification]

Several records by Van der Hammen (1952), RMNH (sub *Scheloribates confundatus*) and Zaitsev & Berg (2001); Hackfort, Vorden, 1959, Minderman (1981), in *Mnium* on border of a ditch, Beekbergen, Elsbosweg, 14.VII.1991, HS.

Liebstadia Oudemans, 1906**306* longior (Berlese, 1908)**

Protoribates badensis Sellnick, 1928

Waterop (LI), VI.1986; Wrakelberg (LI), VI.1986, RMNH (indicated with cf.).

307 similis (Michael, 1888)

Oribatula anomala Hull, 1916

Protoribates silesius Sellnick, 1925

Oribates pallidula; Oudemans 1896 [misidentification]

Several records by Van der Hammen (1952), Zaitsev & Berg (2001) and RMNH; in *Rhytidia delphus squarrosus* on sand dunes, Horsmeertjes, Texel, 6.II.1993, hs.

- Scheloribates Berlese, 1908**
- 308 *laevigatus* (C.L. Koch, 1835)
Oribata lucasi Nicolet, 1855
Oribates fuscomaculata; Oudemans 1896
[misidentification]
Oribates michaeli Hull, 1914
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001), HS, ALT and RMNH.
- 309 *latipes* (C.L. Koch, 1844)
Several records by Van der Hammen (1952)
and RMNH; among litter, Texel, Mokbaai,
6.II.1993, HS.
- 310 *pallidulus* (C.L. Koch, 1840)
Several records by Van der Hammen (1952)
and RMNH; moss under *Sambucus*, Texel,
Horsmeertjes, 6.II.1993, HS.
- Siculobata Grandjean, 1953**
- 311 *leontycha* (Berlese, 1910)
Oisterwijk in Oudemans collection, Van
der Hammen (1952); Ameland, Nesserbos,
28.VI.1972, RMNH.
- Topobates Grandjean, 1958**
- 312 *holsaticus* Weigmann, 1969
Several records from the sea shore in the prov-
ince of Zeeland (Polderman & de Vries 1978);
Noordpolder (GR), 15.XII.1975; Texel, Het
Noorden, 14.XII.1975, RMNH.
- ORIBATULIDAE**
- Oribatula Berlese, 1895**
- 313 *tibialis* (Nicolet, 1855)
Zygoribatula exilis; Buitendijk 1945 [misiden-
tification]
Zygoribatula venustus; Buitendijk 1945 [mis-
identification]
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; Hackfort,
Vorden, 1959, Minderman (1981); pine litter,
Ysselstein (LI), 28.III.1993, HS.
- Phauloppia Berlese, 1908**
- 314 *lucorum* (C.L. Koch, 1841)
Oppia conformis Berlese, 1895
Eremaeus schneideri Oudemans, 1900
Eremaeus conjunctus Oudemans, 1902
Oribata geniculatus Willmann, 1931
Several records by Van der Hammen (1952)
and RMNH; moss on roof tiles, Hoog Soeren,
4.VIII.1989, HS.
- Zygoribatula Berlese, 1916**
- 315* *cognata* (Oudemans, 1902)
Zygoribatula diversa Mihelcic, 1956
Mook, De Diepen, 23.VIII.1989, ALT (speci-
mens not stored).
- 316 *excavata* (Berlese, 1916)
Zygoribatula terricola Van der Hammen,
1952
Zygoribatula cognata; Sellnick 1960 [misiden-
tification]
Valkeveen; Zeeburg; Maasland (typelocality),
12.VIII.1949; Lekkerkerk, 29.X.1949, Van der
Hammen (1952), RMNH, Oudemans collection.
- 317 *exilis* (Nicolet, 1855)
Several records by Van der Hammen (1952),
Zaitsev & Berg (2001) and RMNH; litter
between *Equisetum hyemale*,
Bekkendelle, Winterswijk, 9.IV.1993, HS.
- 318 *frisiae* (Oudemans, 1900)
Several records by Van der Hammen (1952)
and RMNH.

ACKNOWLEDGEMENTS

The authors are indebted to D.A. Krivolutsky for checking the identification of some of the species. Roy Kleukers has been so kind to provide the gross data of the Museum Naturalis Leiden to check and incorporate in this study. Special thanks to Nico M. van Straalen and H.R. Zoomer for some valuable comments concerning this manuscript. The research was done under the financial support of the project 'Biodiversity of soil cryptobiota' awarded by the Institute of Ecological Science, Vrije Universiteit, Amsterdam.

REFERENCES

- Balogh J. & P. Balogh 1992. Oribatid mites genera of the world 1-2. – Hungarian Natural History Museum, Budapest.
- Coddington J.A., G. Giribet, M.S. Harvey, L. Prendini & D.E. Walter 2004. Arachnida. – In: Cracraft, J. & M.J. Donahue (Eds.), Assembling the Tree of Life. Oxford University Press: 296-318.
- Hammen, L. van der 1952. The oribatei (Acari) of the Netherlands. – *Zoölogische Verhandelingen* 17: 1-139.
- International Commission for Zoological Nomenclature 1999. International Code of Zoological Nomenclature, adopted by the xx General Assembly of the International Union of Biological Sciences. – Web edition, Int. Trust Zool. Nomen., London, www.iczn.org/iczn/index.jsp.
- Krivolutsky D.A. (ed.) 1995. Oribatid mites: morphology, development, phylogeny, ecology, methods of study, model species *Nothrus palustris* C.L. Koch, 1839. – Nauka Publishers, Moscow. [in Russian]
- Luxton, M. 1996. Oribatid mites of the British Isles: A check-list and notes on biogeography (Acari: Oribatida). – *Journal for Natural History* 30: 803-822.
- Minderman, G. 1981. Bodemkundige en bosecologische gegevens van het eikenhakhoutbosje Hackfort in de gemeente Vorden. – Rijksinstituut voor Natuurbeheer, Arnhem. [4 delen]
- Niedbala, W. 2004. Fauna Europaea: Acariformes, Oribatida. – Fauna Europaea version 1.1, www.faunaeur.org.
- Noordijk, J. & M. Berg 2001. De corticale fauna van platanen: 1. Arachniden (Arachnida: Araneae, Pseudoscorpiones, Acari). – Nederlandse Faunistische Mededelingen 15: 13-31.
- Oudemans A.C. 1896. List of Dutch Acari Latr. 1. Oribatei. Dug., with synonymous notes and other remarks. – *Tijdschrift voor Entomologie* 39: 53-65.
- Oudemans A.C. 1902. New list of Dutch Acari, 2. With remarks on known and descriptions of a new sub-family, new genera and species. – *Tijdschrift voor Entomologie* 45: 1-52.
- Polderman, P.J.G. 1974. The Oribatida (Acari) of saline areas in the western part of the Dutch Wadden Sea. – *Netherlands Journal of Sea Research* 8: 49-72.
- Polderman, P.J.G. 1977. *Scutovertex pilosetus* nov. spec. from the Netherlands (Acarida, Oribatida). – *Entomologische Berichten* 37: 129-132.
- Polderman, P.J.G. & R. de Vries 1978. Notities over vier soorten mosmijten (Acarida: Oribatida), nieuw voor de Nederlandse fauna. – *Bijdragen Faunistiek Nederland* 5: 20-27.
- Schelvis, J. 1992. Mites and archaeozoology. General methods; applications to dutch sites. – Groningen University. [PhD-thesis]
- Subias, L.S. 2004. Listado sistemático, sinónimico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del mundo (1758-2002). – *Graellsia* 60 (número extraordinario): 3-305.
- Walter D.E. & H.C. Proctor 1999. Mites: Ecology, Evolution and Behaviour. – University of NSW Press, Sydney & CABI, Wallingford.
- Weigmann, G. 2006. Hornmilben (Oribatida). – Goecke & Evers, Keltern. [Die Tierwelt Deutschlands 76]
- Zaitsev, A & M.P. Berg 2001. Some preliminary distribution maps of oribatid mites of the Netherlands (Acari, Oribatida). – Nederlandse Faunistische Mededelingen 14: 79-102.

SUMMARY

Een nieuwe naamlijst van de mosmijten van Nederland (Acari: Oribatida)

In dit artikel wordt een nieuwe naamlijst gepresenteerd van de mosmijten van Nederland.

Voor dit eerste eerste overzicht sinds Van der Hammen (1952) werd de literatuur en collecties bestudeerd. Het aantal bekende soorten werd bijna verdubbeld, tot 318 soorten, waarvan er hier 100 voor het eerst gepresenteerd worden.

H. Siepel

ALTERRA, Wageningen UR and Radboud University Nijmegen.

Center for Ecosystem Studies Animal ecology and physiology group.

P.O. Box 47

6700 AA Wageningen

Henk.Siepel@wur.nl

A.S. Zaitsev

Vrije Universiteit

Institute of Ecological Science, Department Animal Ecology

De Boelelaan 1087

1081 HV Amsterdam

Zaitsev@bio.vu.nl

M.P. Berg

Vrije Universiteit

Institute of Ecological Science, Department Animal Ecology

De Boelelaan 1087

1081 HV Amsterdam

berg@bio.vu.nl

