

Notes on *Pachliopta* species: *Pachliopta phegeus* (Hopffer, 1866) (Lepidoptera: Papilionidae)

Jan J.M. Moonen

KEY WORDS
buraki, homonym, *leytensis*

Entomologische Berichten 75 (4): 132-136

Some confusion has existed about the relationship between *Pachliopta phegeus* and *P. phegeus leytensis* Murayama (1978), which was considered a separate species by some authors. Both taxa occur in the Philippines, and *P. phegeus* is always present where *P. phegeus leytensis* occurs. Nowadays both are seen as forms of a single species. Formerly, *Pachliopta phegeus* was placed in the genus *Papilio*. However, there also is a misunderstanding about the nomenclature of the name *Papilio phegeus* Hopffer. That name and *Papilio phegea* Borkhausen were incorrectly declared homonyms (i.e., the same name ascribed to two species), because both *phegeus* and *phegea* were thought to be adjectives, thus male and female versions of the same word. A new name seemed to be necessary for the newer of the two names, which was *Papilio phegeus*, later *Pachliopta phegeus*, which became *Pachliopta buraki* Koçak. After that, the (older) form name *leytensis* was used in place of *buraki*. However, the names *phegeus* and *phegea* are both proper nouns and thus cannot be homonyms. This means *phegeus* is still valid and the species should be named *Pachliopta phegeus*, with the nominate form f. *phegeus* and the later described form f. *leytensis*.

Introduction

For a long time, *Pachliopta phegeus* Hopffer, 1866 was a quite unknown species, that often only was mentioned in the literature without further information (see list below). More than a century after its discovery, a new subspecies *P. phegeus leytensis* was described by Murayama (1978). However, the taxonomic relationship of this taxon with *Pachliopta phegeus* was unclear. Page & Treadaway (1997, 2003, 2004) considered both taxa as forms of a single species.

Since 1983, *Papilio phegeus* Hopffer, 1866 is considered to be a primary homonym (the same name originally used for two species) of *Papilio phegea* Borkhausen, 1788 (Koçak 1983, Bridges 1988, Page & Treadaway 1997, 2003, 2004). This had strange consequences for the nomenclature of the species, as explained below. All figured specimens are from the collection of the author, now in Naturalis Biodiversity Centre, Leiden (RMNH).

Historical review of *Papilio phegeus*

Hopffer 1866	<i>Papilio phegeus</i> Hopffer
Semper 1886-1892	<i>Papilio</i> (Menelaides) <i>phegeus</i> Hopffer
Rothschild 1895	<i>Papilio phegeus</i> Hopffer
Jordan 1909	<i>Papilio phegeus</i> Hopffer
Bryk 1930	<i>Papilio phegeus</i> Hopffer
Ford 1944	introduces <i>Atrophaneura</i> [for current <i>Parides</i> , <i>Pharmacophagus</i> , <i>Atrophaneura</i> and <i>Pachliopta</i>]
Munroe 1961	<i>Pachlioptera phegeus</i> Hopffer (misspelling of <i>Pachliopta</i>)
Hiura & Alagar 1971	<i>Pachliopta phegeus</i> Hopffer
Smart 1976	<i>Pachliopta phegeus</i> Hopffer
Murayama 1978	<i>Pachliopta phegeus leytensis</i> new subspecies
Igarashi 1979	<i>Pachliopta phegeus</i> Hopffer
Hancock 1980	<i>Atrophaneura</i> (<i>Atrophaneura</i>) <i>phegeus</i> Hopffer
Tsukada & Nishiyama 1980	<i>Pachliopta phegeus</i> Hopffer
	<i>Pachliopta leytensis</i> Murayama (with list of arguments for species status)
D'Abrera 1982	<i>Pachliopta phegeus</i> Hopffer (? = <i>leytensis</i> Murayama)
Hancock 1983	<i>Atrophaneura</i> (<i>Atrophaneura</i>) <i>phegeus</i>
Koçak 1983	<i>Pachliopta buraki</i> [as replacement for <i>phegeus</i> Hopffer, 1866 nec <i>phegea</i> Borkhausen, 1788]

Moonen 1984	<i>Pachliopta phegeus leytenis</i> Murayama (status uncertain.)
Collins & Morris 1985	<i>Atrophaneura</i> (<i>Pachliopta</i>) <i>phegeus</i> Hopffer (<i>leytenis</i> Murayama synonym)
Hancock 1988	<i>Atrophaneura</i> (<i>Pachliopta</i>) <i>phegeus</i> Hopffer
Bridges 1988 [15.iv.1988]	<i>Atrophaneura</i> (<i>Atrophaneura</i>) <i>leytenis</i> Murayama subspecies <i>buraki</i> Koçak (= <i>phegeus</i> Hopffer)
[in Annotations 28.ix.1988]	<i>Atrophaneura</i> (<i>Pachliopta</i>) <i>leytenis</i> Murayama subspecies <i>buraki</i> Koçak (= <i>phegeus</i> Hopffer)
Treadaway 1989	<i>Pachliopta phegeus phegeus</i> Hopffer f. <i>leytenis</i> Murayama
Parsons 1996	<i>Atrophaneura</i> (<i>Pachliopta</i>) [polydorus-group]
Page & Treadaway 1997	<i>Pachliopta leytenis</i> Murayama (syn. <i>P. buraki</i> Koçak = <i>P. phegeus</i> Hopffer) f. <i>leytenis</i> f. <i>buraki</i>
Parsons 1998	<i>Atrophaneura</i> [polydorus (Linnaeus)]
Page & Treadaway 2003 Plates idem as 2004 (Pl. 25, 26)	
Page & Treadaway 2004 Text	<i>Pachliopta leytenis</i> Murayama (syn. <i>P. buraki</i> Koçak = <i>P. phegeus</i> Hopffer) <i>P. leytenis</i> f. <i>leytenis</i> <i>P. leytenis</i> f. <i>buraki</i>
Häuser et al. 2005	<i>Pachliopta leytenis</i> Murayama
Racheli & Cotton 2010	<i>Pachliopta</i> checklist, in which <i>phegeus/leytenis</i> was unintentionally omitted (Cotton pers. com.).

Not a primary homonym

Koçak (1983) stated that *Papilio phegeus* Hopffer, 1866 is a primary homonym of *Papilio phegea* Borkhausen, 1788, a species currently known as *Protoerebia afra* (Fabricius), Nymphalidae, Satyrinae. He based his conclusion on the assumption that *phegeus* and *phegea* are both adjectives. If this was the case, *phegea* would be the female form of *phegeus* (incorrectly used because the gender of *Papilio* is male), and both words would in fact be the same. Consequently Koçak replaced the newer name, Hopffer's *phegeus*, with *buraki*. This is, however, incorrect. A thorough examination of several Greek and Latin dictionaries revealed that both are proper nouns. *Phegeus* (Φηγεύς) was a king of the fortified city of Psophis in Northwest Arcadia (in the centre of the Greek Peloponnese). *Phegea* (Φηγέα) is the earlier

name of Psophis. A king and a city are clearly not the same, which means that *Papilio phegeus* Hopffer cannot be a homonym of *Papilio phegea* Borkhausen, in the same way that *Papilio helenus* Linnaeus, 1758 and *Papilio helenia* Linnaeus, 1758 (now *Troides helenia*) are not homonyms. Thus there is no need to replace *phegeus* by *buraki*.

There has been some disagreement in the literature as to which genus the taxon belongs in, either *Pachliopta* or *Atrophaneura* (see the list). This paper does not deal with this higher level taxonomic problem. Here, Racheli & Cotton (2010) and other authors are followed in placing *phegeus* in the genus *Pachliopta*.



1. ♂ *Pachliopta phegeus* f. *phegeus* Hopffer. Philippines: Southern Leyte: Sta. Cruz, St. Bernard, 4.iii.1979. Left: upperside, right: underside (length forewing 47 mm). Photo: Jan Moonen
1. ♂ *Pachliopta phegeus* f. *phegeus* Hopffer. Filippijnen: Southern Leyte: Sta. Cruz, St. Bernard, 4.iii.1979. Links: bovenzijde, rechts: onderzijde (lengte voorvleugel 47 mm).



2. ♂ *Pachliopta phegeus* f. *leytensis* Murayama. Philippines: Southern Leyte: Catmon St. Bernard, 22.iv.1979. Left: upperside, right: underside (length forewing 51 mm). Photo: Jan Moonen
2. ♂ *Pachliopta phegeus* f. *leytensis* Murayama. Filippijnen: Southern: Leyte, Catmon St. Bernard, 22.iv.1979. Links: bovenzijde, rechts: onderzijde (lengte voorvleugel 51 mm).




3. ♀ *Pachliopta phegeus* f. *phegeus* (Hopffer). Philippines: Bohol, 20.viii.1990. Left: upperside, right: underside (length forewing 54 mm). Photo: Jan Moonen
3. ♀ *Pachliopta phegeus* f. *phegeus* (Hopffer). Filippijnen: Bohol, 20.viii.1990. Links: bovenzijde, rechts: onderzijde (lengte voorvleugel 47 mm).

Variation within *Pachliopta phegeus* (Hopffer)

In 1978, Murayama described a new subspecies for *P. phegeus*: *leytensis* (figure 1-4). Tsukada & Nishiyama (1980) mentioned *Pachliopta leytensis* as a species distinct from *Pachliopta phegeus* based on partial sympatry and the apparent absence of intermediate specimens, although there are no differences in the male genitalia. Authors who followed the unjustified action of Koçak (1983), concluded that *leytensis* is the oldest available

name for *Pachliopta phegeus*, so its name should be changed to *Pachliopta leytensis* instead of *Pachliopta buraki*. Some considered *buraki* to be a form or subspecies of it, or a species of its own, leading to a confusing taxonomic and nomenclatural situation. Page & Treadaway (2004), still following Koçak (1983) and mentioning that intermediate specimens were known known, described the taxonomy of the species as follows:

<i>Pachliopta leytensis</i> Murayama, 1978		
f. <i>leytensis</i> Murayama, 1978	[S. Leyte, E. Mindanao, Samar, Siargo]	
f. <i>buraki</i> Koçak, 1983	[= <i>phegeus</i> Hopffer, 1866 nec <i>phegea</i> Borkhausen, 1788] [Bohol, Cebu, Dinagat, S. Leyte, S.-C. & E. Mindanao, Panaon, Samar, Siargo]	



4. ♀ *Pachliopta phegeus* f. *leytensis* Murayama. Philippines: Southern Leyte: Hinunangan, 4.ii.1992. Left: upperside, right: underside (length forewing 50 mm). Photo: Jan Moonen
4. ♀ *Pachliopta phegeus* f. *leytensis* Murayama. Filippijnen: Southern Leyte: Hinunangan, 4.ii.1992. Links: bovenzijde, rechts: onderzijde (lengte voorvleugel 47 mm).

Taking the foregoing into account, the taxonomy/nomenclature has to be corrected to its original situation:

Pachliopta phegeus (Hoppfer, 1866) [status revised]
f. *phegeus* (Hoppfer, 1866) (= *buraki* Koçak, 1983) [status revised]
f. *leytensis* Murayama, 1978 [status revised]

Acknowledgements

I thank Rienk de Jong for his critical comments. Further I thank the Uyttenboogaart-Eliassen Foundation for the financial support

that makes it possible to travel frequently between Cadix en Keer and Leiden, Naturalis.

Literature

- Bridges CA 1988. Catalogue of Papilionidae & Pieridae (Lep.: Rhop.). Bridges.
- Bryk F 1930. Papilionidae II (*Papilio*). In: Lepidopterorum Catalogus Pars 37 (Strand E ed): 57-510. Junk.
- Collins NM & Morris MG 1985. Threatened swallowtail butterflies of the World. The IUCN Red Data Book. IUCN.
- D'Abreu B 1982. Butterflies of the Oriental Region. Part I Papilionidae, Pieridae & Danaidae. Hill House.
- Ford EB 1944. Studies on the chemistry of pigments in the Lepidoptera, with reference to their bearings on systematics. 4. The classification of the Papilionidae. Proceedings of the Royal Entomological Society, London A 19: 92-106.
- Hancock DL 1980. The status of the genera *Atrophaneura* Reakirt and *Pachliopta* Reakirt (Lepidoptera: Papilionidae). Australian Entomological Magazine 7(2): 27-32.
- Hancock DL 1983. Classification of the Papilionidae (Lep.): a phylogenetic approach. *Smithersia* 2: 1-48.
- Hancock DL 1988. A revised classification of the genus *Atrophaneura* Reakirt (Lepidoptera: Papilionidae). Australian Entomological Magazine 15: 7-16.
- Häuser CL, De Jong R, Lamas G, Robbins RK, Smith C & Vane-Wright RI 2005. Papilionidae – revised GloBIS/GART species checklist (2nd draft). Available at: www.insects-online.de/frames/papilio.htm [accessed January 2012]
- Hiura I & Alagar RE 1971. Studies on the Philippine butterflies chiefly collected by the co-operative survey by the Osaka Museum of Nat. Hist. and the National Museum of the Philippines, 1969. Part 1: Papilionidae. Bulletin of the Osaka Museum of Natural History 24: 29-44.
- Hoppfer CH 1866. Neue Arten der Gattung *Papilio* im Berliner Museum. Entomologische Zeitung, herausgegeben von dem entomologischen Vereine zu Stettin 27(1-3): 22-32.
- Igarashi S 1979. Papilionidae and their early stages, vol 1 & vol 2 [in Japanese]. Kodansha.
- Jordan K 1908-1910. Papilionidae, *Papilio* bis *Armandia*. In: Seitz: Großschmetterlinge der Erde, die Indo-Australische Tagfalter IX: 11-109.
- Koçak AO 1983. More notes on the homonymy of the specific names of Lepidoptera (continued). *Priamus* 2: 164-166.
- Moonen JJM 1984. Notes on Eastern Papilionidae. *Papilio International* 1: 47-50.
- Munroe E 1961. The Classification of the Papilionidae (Lep.). The Canadian Entomologist, suppl. 17: 1-51.
- Murayama S 1978. On some species of Rhopalocera from Southeast Asia with descriptions of a new species and a new subspecies. *Tyô To Ga* 29: 153-158.
- Page MGP & Treadaway CG 1997. Revision of *Pachliopta phlegon* (C. & R. Felder, 1864), *P. buraki* Koçak, 1983 and *P. leytensis* Murayama, 1978 (Lep., Pap.). Nachrichten des entomologischen Vereins, suppl. 17: 377-392.
- Page MGP & Treadaway CG 2003. Papilionidae of the Philippine Islands. In: Butterflies of the World. part 17 Papilionidae IX (Bauer E & Frankenbach T eds). Goecke & Evers.
- Page MGP & Treadaway CG 2004. Papilionidae of the Philippine Islands. In: Butterflies of the World. Supplement 8 (Bauer E & Frankenbach T eds). Goecke & Evers.
- Parsons MJ 1996. Gondwanan evolution of the Troidine Swallowtails (Lep.: Pap.): Cladistic reappraisals using mainly immature stage characters, with focus on the birdwings *Ornithoptera* Boisduval. Bulletin of the Kitakyushu Museum of Natural History 15: 43-118.
- Parsons M 1998. The Butterflies of Papua New Guinea, their Systematics and Biology. Academic Press.
- Racheli T & Cotton AM 2010. Papilionidae


- part II. In: Guide to the Butterflies of the Palaearctic Region (Bozano GC ed). Omnes Artes.
- Rothschild W 1895. A Revision of the Papilios of the eastern Hemisphere, exclusive of Africa. Novitates zoologicae II(3): 167-463.
- Semper G 1886-1892. Die Schmetterlinge der Philippinischen Inseln. Beitrag zur Indo-Malayischen Lepidopteren-Fauna. 1 (Die Tagfalter). Reisen im Archipen der Philippinen (2) 5. Wiesbaden 1.
- Smart P 1976. The illustrated Encyclopedia of the Butterfly World. Salamander Books.
- Treadaway CG 1989. A check list of the Philippine Papilionidae. Papilio International 5: 375-381.
- Tsukada E & Nishiyama Y 1980. Papilionidae. In: Butterflies of the South East Asian Islands (Tsukada E ed). Plapac Co.

Accepted: 4 February 2015

Samenvatting

Notities over *Pachliopta*: *Pachliopta phegeus* (Hopffer, 1866) (Lepidoptera: Papilionidae)

Er was enige verwarring over de relatie tussen *Pachliopta phegeus* en de pas in 1978 beschreven *Pachliopta phegeus leytenensis* Murayama, die door sommige auteurs als aparte soort werd beschouwd. Beide vlinders komen in de Filippijnen voor. Waar *leytenensis* vliegt, komt ook altijd *phegeus* voor. Nu beschouwt men beide als vormen (forma's) van één soort. Echter, er bestond ook een misverstand over de nomenclatuur van de soort waar beide vormen onder vallen. *Pachliopta phegeus* was oorspronkelijk beschreven in het genus *Papilio*. Onterecht waren *Papilio phegeus* (Hopffer) en *Papilio phegea* Borkhausen tot homonymen (dezelfde naam voor twee soorten) verklaard, ervan uitgaande dat *phegeus* en *phegea* de mannelijke en vrouwelijke vorm van hetzelfde bijvoeglijk naamwoord waren. Een nieuwe naam, *buraki* Koçak, leek nodig te zijn voor de jongste van de twee namen: *P. phegeus* werd *P. buraki*. Intussen was er een ondersoort van *P. phegeus* beschreven: *P. phegeus leytenensis*. Daarmee werd *leytenensis* de oudst beschikbare naam voor de soort: *P. buraki* werd van toen af *P. leytenensis*. De namen *phegeus* en *phegea* zijn echter beide zelfstandige naamwoorden en zijn dus geen homonymen van elkaar. Dat wil zeggen dat *phegeus* zijn geldigheid terug heeft en de soort weer *Pachliopta phegeus* heet, met de nominaatvorm f. *phegeus* en de later beschreven vorm f. *leytenensis*.



Jan J.M. Moonen
Willem Alexanderstraat 4
6267 AR Cadier en Keer
The Netherlands
jan@moonen.net