

A new species of living scallop of the genus *Anguipecten* (Bivalvia, Pectinidae)
from the tropical Indo-Pacific

HENK H. DIJKSTRA

c/o Zoological Museum, University of Amsterdam, P.O. Box 94766, NL 1090 GT Amsterdam, The
Netherlands; h.h.dijkstra@wxs.nl

Anguipecten pacificus spec. nov. is described from the tropical Indo-Pacific.

Key words: Bivalvia, Pectinidae, *Anguipecten*, new species, Marquesas Islands, Indonesia.

INTRODUCTION

During the Indonesian-French Karubar cruise to eastern Indonesia a new scallop was found near the Kai Islands, which was provisionally identified by Dijkstra & Kastoro (1997: 279, figs 156-159), based on a few dead specimens from the upper bathyal depths. Subsequently, more material was collected by the French Musorstom 9 cruise to the Marquesas Islands (French Polynesia), sufficient to describe it herein. All specimens (empty shells) were deposited in MNHN, with voucher specimens in the private pectinoid reference collection of the author.

Acronyms and abbreviations: HD, H.H. Dijkstra colln, Sneek, The Netherlands; MNHN, Muséum national d'Histoire naturelle, Paris, France. For shell characters: D, diameter (thickness of both valves); H, height (dorsal-ventral); lv, left valve (upper valve); pr, paired (articulated) valves; rv, right valve (lower valve); W, width (anterior-posterior).

SYSTEMATIC PART

Subfamily Pectininae Wilkes, 1810

Tribe Decatopectinini Waller, 1986

Anguipecten Dall, Bartsch & Rehder, 1938

Anguipecten Dall, Bartsch & Rehder, 1938: 92. Type species (by original designation): *Anguipecten gregoryi* Dall, Bartsch & Rehder, 1938 [junior synonym of *Pecten lamberti* Souverbie, in Souverbie & Montrouzier, 1874], Miocene to Recent, tropical Indo-Pacific (Hayami, 1989: 15).

Anguipecten pacificus spec. nov. (figs 1-4)

Anguipecten spec. cf. *picturatus* Dijkstra, 1995; Dijkstra & Kastoro, 1997: 279, figs 156-159.

Material examined (in MNHN, unless stated otherwise). — French Polynesia, Marquesas Islands (Musorstom 9 cruise; Bouchet, Dayrat & Richer de Forges leg.): Stn DW 1274, Eiao Island, 07°54.6'S

140°40.1'W, 100-120 m, 3 paratypes (1 lv, 2 rv); Stn DW 1281, Motu One, 07°48'S 140°21'W, 450-455 m, holotype (lv), 2 paratypes (rv); Stn DW 1287, Eiao Island, 07°54'S 140°40'W, 163-245 m, 3 paratypes, one of which in HD (lv).

Indonesia (Karubar cruise; Bouchet, Kastoro & Métivier leg.): Stn DW 30, Kai Islands, 05°39'S 132°56'E, 111-118 m, 4 specimens, not considered paratypes (2 lv, 2 rv).

Description. — Shell up to c. 60 mm high, solid, compressed, oblong, nearly equivalve and equilateral, right valve slightly more convex than left valve, auricles short and subequal, umbonal angle c. 85°, colour creamy, maculated with reddish dots and streaks, right valve somewhat paler.

Both valves radially undulated and sculptured with six prominent lirae; laterally an additional but weakly developed ridge. Interstices somewhat narrower than lirae. Secondary radial sculpture of small lirae (on right valve weaker) on the lirae and interstices, commence on the central part of the disc (c. 10 per lira) and extend to the ventral margin. Microsculpture consists of delicate and closely spaced commarginal lamellae throughout (c. 8-10 per mm in the central part). Anterior auricles very weakly sculptured with c. 10 radial riblets, posterior auricles nearly smooth. Inner surface plicated, more prominently so near periphery. Resilifer triangular. Byssal notch, fasciole and ctenolium lacking. Cardinal crura rather broad with closely and vertically set small ridges.


Measurements of type series: holotype H 40.2 mm, W 35.8 mm; largest paratype H 60.5 mm.

Distribution. — Eastern Indonesia (Kai Islands) and French Polynesia (Marquesas Islands); only dead taken bathyally (100-455 m), probably living sublittorally according to other data of *Anguipecten* species from the tropical Indo-Pacific (Dijkstra, unpubl. data).

Comparison. — The specimens of *A. pacificus* from eastern Indonesia are somewhat atypical and slightly differ from the type material by having somewhat larger auricles with a shallow byssal notch on the right valve, which is lacking in typical material. Therefore these specimens are excluded from the type series. *Anguipecten superbus* (G.B. Sowerby II, 1842), known from the Indo-West Pacific, is very similar in size, shape, colour, and microsculpture, but differs from *A. pacificus* in having many more (18-23, *A. pacificus*: 6-7) radial lirae and a far less prominent secondary sculpture. *Anguipecten lamberti* (Souverbie, in Souverbie & Montrouzier, 1874), known from New Caledonia and Hawaiian Islands, differs in having numerous (c. 30) very compressed radial lirae and a weak secondary radial sculpture. Other morphological characters are similar to those of *A. pacificus* and *A. superbus* (size, shape, colour, and microsculpture).

Anguipecten picturatus Dijkstra, 1995, also known from the Indo-West Pacific, differs from *A. pacificus* in size (up to c. 45 mm in height), in colour (more bright), in having more radial lirae (12-15), a coarser secondary radial sculpture, and a commarginal microsculpture in the late growth stage.

Etymology. — From the Pacific Ocean (Lat. *pacificus*, adjective = pacific, peaceable).



Figs 1-9. *Anguipecten* spec. Figs 1-4. *A. pacificus* spec. nov., Marquesas Islands, Motu One, Musorstom 9, Stn DW 1281, 07°48'S 140°21'W, depth 450-455 m, dead, N/O "Alis", 07.ix.1997. 1-2, holotype (H 40.2 mm, W 35.8 mm); 1, lv, exterior; 2, lv, interior. 3-4, paratype (H 40.0 mm, W 35.8 mm); 3, rv, exterior; 4, rv, interior (MNHN, not registered). Figs 5-7. *A. lamberti* (Souverbie, in Souverbie & Montrouzier, 1874). 5-6, New Caledonia, Boula Ri Pass, outside of reef, 60 m, alive, soft sediments, 1985; pr (H 69.9 mm, W 63.1 mm, D 15.9 mm). 5, lv, exterior; 6, lv, interior (HD 2535); 7, Hawaiian Islands, Oahu, 73 m, alive, under slab, c. 1988; pr (H 36.8 mm, W 35.2 mm, D 6.8 mm), lv and rv, exterior (HD 7237). Fig. 8. *A. picturatus* Dijkstra, 1995, Japan, Okinawa, Horseshoe, depth 40 m, alive, soft sediments, 1985; pr (H 47.9 mm, W 42.8 mm, D 13.9 mm), lv and rv, exterior (HD 1949). Fig. 9. *A. superbus* (Sowerby II, 1842), Solomon Islands, Ontong Java Atoll, 20 m, alive, soft sediments, 1986; pr (H 58.1 mm, W 55.8 mm, D 15.5 mm), lv and rv, exterior (HD 2937).

ACKNOWLEDGEMENTS

I am much indebted to Dr P. Bouchet (Muséum national d'Histoire naturelle, Paris) for submitting this pectinid material for study purposes. Thanks are also due to Mr P. Maestrati (Muséum national d'Histoire naturelle, Paris) and Mr R.G. Moolenbeek (Zoological Museum, University of Amsterdam, Amsterdam) for their technical assistance and cooperation.

REFERENCES

- DALL, W.H., P. BARTSCH & H.A. REHDER, 1938. A manual of the Recent and fossil marine pelecypod mollusks of the Hawaiian Islands. — Bernice P. Bishop Museum Bulletin 153: i-iv, 1-233. Honolulu.
- DIJKSTRA, H.H., 1995. Notes on taxonomy and nomenclature of Pectinidae (Mollusca: Bivalvia). 1. *Anguipecten picturatus* nom. nov. — Basteria 59: 15-19.
- DIJKSTRA, H.H., & W.W. KASTORO, 1997. Mollusca Bivalvia: Pectinoidea (Propeamussiidae and Pectinidae) from eastern Indonesia. In: A. CROSNIER & P. BOUCHET, eds, Résultats des Campagnes MUSORSTOM, vol. 16. — Mémoires du Muséum national d'Histoire naturelle 172: 245-285. Paris.
- HAYAMI, I., 1989. Outlook on the Post-Paleozoic historical biogeography of pectinids in the Western Pacific region. In: I. OHBA, I. HAYAMI & K. MOCHIZUKI, eds, Current aspects of biogeography in West Pacific and East Asian regions: 1-25. Tokyo.
- SOUVERBIE, [S.M.], & R.P. MONTROUZIER, 1874. Descriptions d'espèces nouvelles de l'Archipel calédonien. — Journal de Conchyliologie 22: 187-201.
- SOWERBY, G.B. 2nd, 1842. Monograph of the genera *Pecten* and *Hinnites*. In: Thesaurus Conchyliorum, or figures and descriptions of Recent shells 1 (2): 45-82, pls 12-20. London.
- WALLER, T.R., 1986. A new genus and species of scallop (Bivalvia: Pectinidae) from off Somalia, and the definition of a new tribe Decatopectinini. — The Nautilus 100: 39-46.